



EUROBATS



EUROBATS

Publication Series
No.

4

Achieving and maintaining favourable conservation status for bats requires that their breeding and resting sites – roosts – are protected. In Europe, a high percentage of bat species roost for at least part of each year in buildings. Buildings of cultural heritage importance (e.g. churches, castles, bridges) are often of particular importance for bats. These structures may be protected in their own right leading to conflict between building conservation work and bat conservation.

To examine this issue EUROBATS established a working group to gather information on bat species roosting in overground buildings; the types of roosts used; and the methods employed to protect these roosts, particularly those in buildings of cultural heritage importance.

A questionnaire addressing these topics was circulated to all range states and responses were received from 50 experts from 37 countries. This publication summarises those responses. It also provides practical advice, illustrated by case studies, on ways in which the potential conflicts between the protection of bats and the conservation of heritage buildings can be managed.



Protection of overground roosts for bats

ISBN 978-92-95058-17-0
(printed version)

ISBN 978-92-95058-18-7
(electronic version)

Ferdia Marnell • Primož Presetnik



Marnell, F. & P. Presetnik (2010): Protection of overground roosts for bats (particularly roosts in buildings of cultural heritage importance). EUROBATS Publication Series No. 4 (English version). UNEP/EUROBATS Secretariat, Bonn, Germany, 57 pp.

Produced by	UNEP/EUROBATS
Coordinator	Christine Boye/EUROBATS Secretariat
Editors	Christine Boye, Tine Meyer-Cords
Proofreading	Kate Horn
Design	Karina Waedt

2nd, updated version 2010 — © 2010 Agreement on the Conservation of Populations of European Bats (UNEP/EUROBATS).

This publication may be reproduced in whole or in part and in any form for educational or non-profit purposes without special permission from the copyright holder, provided acknowledgement of the source is made. UNEP/EUROBATS would appreciate receiving a copy of any publication that uses this publication as a source.

No use of this publication may be made for resale or for any other commercial purpose whatsoever without prior permission in writing from UNEP/EUROBATS.

The boundaries shown on the maps in this publication do not imply official endorsement or acceptance by the United Nations.

Copies of this publication are available from the

UNEP/EUROBATS Secretariat
United Nations Campus
Hermann-Ehlers-Str. 10
53113 Bonn, Germany
Tel (+49) 228 815 24 21
Fax (+49) 228 815 24 45
E-mail eurobats@eurobats.org
Web www.eurobats.org

ISBN 978-92-95058-17-0 (printed version)
ISBN 978-92-95058-18-7 (electronic version)

Cover photo: Church in Zavrse, Slovenia.
© P. Presetnik

UNEP promotes environmentally sound practices globally and in its own activities. This publication is printed on chlorine-free, FSC-certified, 60% recycled paper, using environmentally friendly practices. Our distribution policy aims to reduce UNEP's carbon footprint.



Contents

Index of tables	5	
Index of figures	5	
Index of case studies	6	
1	Introduction	7
2	Literature Review	8
3	Use of overground roosts by bats	9
3.1	Bat species present in overground roosts	9
3.2	Geographic pattern of dependence of bat species on overground roost types	12
3.3	Intraspecific variation across the European range	13
4	Roost protection	19
4.1	Legal protection	19
4.2	Physical protection	20
4.3	Education/information	20
4.3.1	Websites	20
4.3.2	Telephone helplines	21
4.3.3	Site notices	21
5	Challenges posed by roosts in domestic dwellings	22
5.1	Disturbance or exclusion	24
6	Focus on buildings of cultural heritage	27
6.1	Protection of cultural heritage	27
6.2	Damage by bats in buildings of cultural heritage	28
6.3	Accommodating bats during renovation/restoration	30
6.3.1	Avoidance	30
6.3.2	Incorporating existing roosts into renovated buildings	33
6.3.2.1	Roost size	33
6.3.2.2	Roost entrances	36
6.3.3	Incorporating new roosts into buildings	37
6.3.4	Barns	40
6.3.5	Bridges	41
6.3.5.1	Bridge survey and mitigation measures	41
6.3.6	Timber treatment, pest control and lead poisoning	43



7	Literature and further reading	44
	Scientific and common names of European bats	48
	Acknowledgements	49
	Annex 1: Questionnaire on dependence of bat species on overground roost types	50
	Annex 2: Summary of questionnaire responses	54
	Annex 3: EUROBATS Resolution 5.7: Guidelines for the Protection of Overground Roosts, with particular reference to roosts in buildings of cultural heritage importance	56
	Annex 4: Summary of good practise for the protection of overground roosts in buildings of cultural heritage importance	57

Index of tables

Table 1.	Percentage of EUROBATS range states where bat species are highly dependent on overground roost types.	10
Table 2.	Bats in houses: common scenarios and possible solutions.	23
Table 3.	Optimum season for works in different types of roosts.	31
Table 4.	Specific roost types and sizes for bat species highly dependent on roosts in buildings.	35

Index of figures

Figure 1.	Percentage of European bat species dependent on different overground roost types.	12
Figure 2.	Percentage of bat species highly dependent on overground roosts in potential cultural heritage buildings (castles, churches, houses and barns combined).	13
Figure 3.	Absolute number and percentage of bat species highly dependent on roosts in castles/fortifications in EUROBATS range states.	14
Figure 4.	Absolute number and percentage of bat species highly dependent on roosts in churches in EUROBATS range states.	15
Figure 5.	Absolute number and percentage of bat species highly dependent on roosts in houses/blocks of flats in EUROBATS range states.	15
Figure 6.	Absolute number and percentage of bat species highly dependent on roosts in barns/stables in EUROBATS range states.	16
Figure 7.	Absolute number and percentage of bat species highly dependent on roosts in bridges in EUROBATS range states.	16
Figure 8.	Absolute number and percentage of bat species highly dependent on roosts in trees in EUROBATS range states.	17
Figure 9.	Estimated dependence of <i>Rhinolophus hipposideros</i> on roosts in churches.	17
Figure 10.	Estimated dependence of <i>Myotis myotis</i> on roosts in potential cultural heritage buildings (castles, churches, houses and barns combined).	18
Figure 11.	Example of bat roost notice <i>in situ</i> , Dublin, Ireland.	21
Figure 12.	Block of flats in Slovenia where a roost of <i>Nyctalus noctula</i> under the exterior cladding only became apparent when renovation work commenced.	22
Figure 13.	Damage to stonework at Penmon Priory, Wales, United Kingdom, due to long-term exposure to bat urine.	28
Figure 14.	Ratková Church, Slovak Republic, clean-up team with bagged bat guano.	30



Figure 15. St Cadoc's Church, Wales, United Kingdom, undergoing essential repairs.	32
Figure 16. Northwest tower of Grad na Goričkem castle, Slovenia, during and at the end of reconstruction in year 2006.	34
Figure 17. Dedicated bat roost entrance, Ireland (also used by lesser horseshoes!).	37
Figure 18. Outbuilding at Glaninchiquin, Ireland, before and after renovation for <i>Rhinolophus hipposideros</i> .	38
Figure 19. Old and new <i>Tadarida</i> roosts, Portugal.	39
Figure 20. Paston Barn, England, United Kingdom.	40
Figure 21. Lisconny Bridge, Ireland, after restoration.	42
Figure 22. Underside of Lisconny Bridge.	42
Figure 23. Château de Trévarez, Brittany, France	43

Index of case studies

Case study 1: Ratková Church, Slovak Republic.	30
Case study 2: St Cadoc's Church, Wales, United Kingdom.	32
Case study 3: Grad na Goričkem, Slovenia.	34
Case study 4: Glaninchiquin, Ireland.	38
Case study 5: Morcegário, Portugal.	39
Case study 6: Paston Barn, England, United Kingdom.	40
Case study 7: Lisconny Bridge, Ireland.	42

1 Introduction

In 2003, the 4th EUROBATS Meeting of Parties (MOP4) asked the Advisory Committee (AC) to gather information on methods used to protect roosts other than underground sites, with roosts in buildings that are part of the cultural heritage as a priority.

An Intersessional Working Group (IWG) was established at AC9 in Vilnius, Lithuania, in 2004 to address this issue. A questionnaire (Annex 1) was circulated to all Parties and Non-Party Range States in December 2004. The questionnaire sought information on the types of overground sites used as roosts, on the administrative and practical protection of roosts and on the interactions between bats and buildings of cultural heritage. Responses were received from the following 37 countries: Albania, Armenia, Austria, Azerbaijan, Belgium, Bosnia and Herzegovina, Bulgaria, Croatia, Czech Republic, Denmark, Estonia, Finland, France, Georgia, Germany, Hungary, Ireland, Italy, Latvia, Liechtenstein, Lithuania, Luxembourg, FYR Macedonia, Montenegro (joint questionnaire with Serbia), the Netherlands, Norway, Poland, Portugal, Romania, Russian Federation, Serbia, Slovakia, Slovenia, Sweden, Switzerland, Ukraine and the United Kingdom.

This publication summarises the results from the overground roost questionnaire, provides an overview of suitable protective measures, and explores the interactions between bats and buildings of cultural heritage importance. Emphasis was put on buildings of cultural heritage importance since it was known that they present important roosting sites for many bat species and the priorities of cultural and natural conservation can and often do collide. A number of case studies are included to illustrate how conflicts between bats and cultural heritage have been successfully resolved in different situations in different parts of Europe.



2 Literature Review

The protection of bats in the man-made environment is an area of active research and the interactions between man and bats attract attention throughout Europe. Most of the published literature, however, examines this issue from the bat conservation angle; publications from the built heritage perspective are harder to come by. A large volume of easily available literature on the protection of overground roosts has been published in the United Kingdom. It is these sources, and in particular *Bat Mitigation Guidelines* (MITCHELL-JONES 2004), *The Bat Workers Manual* (MITCHELL-JONES & MCLEISH 2004), THE NATIONAL TRUST'S *Wildlife and Buildings* (2001) and *The Bats in Churches Project* (SARGENT 1995), which largely inform this document. The reader is also referred to two important German publications (also available in English), which examine the eco-

logical requirements of European bats: *Bat roosts in the Alpine area: Guidelines for the renovation of buildings* (REITER & ZAHN 2006) and *Ecology and Conservation of Bats in Villages and Towns* (SIMON *et al.* 2004). Other published and unpublished materials from Austria, Estonia, France, Germany, Ireland, Italy, Latvia, Lithuania, the Russian Federation, Ukraine and the United Kingdom were also examined in the preparation of this report (see chapter 7).

EUROBATS has already produced an advisory document on underground roosts in its Publication Series: *Protecting and managing underground sites for bats* by MITCHELL-JONES *et al.* (2007). This present report aims to complement that underground roost document and where overlaps occur the reader will be referred to that earlier report.

3 Use of overground roosts by bats

As their metabolic and social requirements vary throughout the year, most bats will use a variety of roosts of different types. Some species are particularly closely associated with tree roosts, but the majority use a range of roosts, which includes trees, buildings and underground sites.

Man-made overground structures regularly used by bats across Europe include bridges, castles, churches, houses, blocks of flats, barns and stables. Some species have come to rely on such structures, *e.g.* *Eptesicus* and *Pipistrellus* species usually roost in buildings; *Myotis daubentonii* is, in some countries, particularly associated with bridges and will form roosts in suitable cracks in both old and new structures; *Myotis myotis* can be found roosting in churches over much of its range in the northern part of Europe; and *Plecotus* species have come to rely more and more on man-made roost sites in some countries due to the successive loss of suitable natural habitat.

Bats can be found in buildings all year round. In late spring, maternity roosts are formed in the roofs of buildings to take advantage of the heat provided by the sun. This is because breeding females seek warm areas during this phase in their life-cycle to minimise the energy cost of maintaining a high body temperature. Some species such as *Pipistrellus* spp. show a clear preference for confined roost sites such as soffit boxes, eaves or under hanging tiles, whereas others such as the *Rhinolophus* spp. are more typically associated with open roof

voids that they can fly into. There are many exceptions and many species have been recorded from a wide variety of situations. In winter, bats of most species have been recorded hibernating in various parts of buildings such as inside cavity walls, around window frames, under ridge tiles and in cooler areas with stable temperatures such as cellars and basements. These latter are covered by the EUROBATS report on underground roosts (MITCHELL-JONES *et al.* 2007) and are not considered further here.

3.1 Bat species present in overground roosts

In 2004, a EUROBATS questionnaire asked national experts to estimate the dependence of individual bat species on specific roost types as high; medium; low; not important; not known; or present (when no detailed information was available). A database of the experts' responses can be found at http://www.eurobats.org/publications/publication_series/overground_database.htm. Table 1 provides a summary of the responses; a more detailed breakdown can be found in Annex 2. An analysis was then conducted of the dependence of bats on different overground roost types in different countries. The main roost types identified were castles/fortifications, churches, houses/blocks of flats, barns/stables, bridges and trees.

A number of caveats should be borne in mind when examining the data:

- i) For a large proportion of bat species the degree of dependence on specific

- roost types in specific countries is unknown (see Annex 2).
- ii) Dependence of particular bat species can vary greatly in different regions of the same country, but each particular country was treated as one region and this produced some confusing results, particularly in larger countries.
 - iii) It is not clear if all the answers dealing with castles/fortifications are only referring to overground roost types; some may include underground habitats (cellars, basements etc.).
 - iv) For the analysis and presentation of results the higher dependence has been taken in cases when answers provided intermediate estimates (e.g. high/medium dependence), or were unclear (e.g. high dependence?). When it was indicated that a species is just “present” in a particular roost type we treated this as “unknown” status of dependence.

Results of analysis reveals that across different EUROBATS range countries at least 33 bat species (73% of known species in the EUROBATS zone) are considered to have high or medium dependence on roosts in castles and fortifications; 32 species (71%) on roosts in churches, and houses or blocks of flats; 27 species (60%) on roosts in barns or stables; and 23 species (51%) on roosts in bridges (Figure 1). The percentage of bat species dependant on roosts in trees (Figure 1) is underestimated, because roosts of tree dwelling bats are unknown in many countries (Annex 3).

If we consider castles, churches, houses and barns (Figure 1, Table 1, Annex 2) as probable buildings of cultural importance, we can estimate that the conservation of approximately 75% of bat species across EUROBATS range states rely for at least part of their life cycle on roosts in buildings of cultural heritage importance.

Table 1. Percentage of EUROBATS range states where bat species are highly dependent on overground roost types.

Species	Overground roost type					
	Castle/fortification	Church	House/block of flats	Barn/stable	Bridge	Tree
<i>Barbastella barbastellus</i>	+		+	+		++
<i>Barbastella leucomelas</i>						
<i>Eptesicus bottae</i>			++			++
<i>Eptesicus nilssonii</i>	+	+	++			+
<i>Eptesicus serotinus</i>	+	++	+++	+		
<i>Hypsugo savii</i>			+	+		
<i>Miniopterus schreibersii</i>	+	+				
<i>Myotis alcathoe</i>						++
<i>Myotis aurascens</i>			+	+	+	
<i>Myotis bechsteini</i>						++++
<i>Myotis blythii</i>	+	++	+	+	+	
<i>Myotis brandtii</i>	+		++			++

Table 1 (cont.)

Species	Overground roost type					
	Castle/fortification	Church	House/block of flats	Barn/stable	Bridge	Tree
<i>Myotis capaccinii</i>					+	
<i>Myotis dasycneme</i>	+	++	+++	+		+
<i>Myotis daubentonii</i>	+	+	+		+	++++
<i>Myotis emarginatus</i>	++	++	+	+		
<i>Myotis hajastanicus</i>						
<i>Myotis myotis</i>	++	+++	+			
<i>Myotis mystacinus</i>	+	+	++	+	+	++
<i>Myotis nattereri</i>	+	+	+	+	+	++
<i>Myotis nipalensis</i>						
<i>Myotis punicus</i>						
<i>Myotis schaubi</i>						
<i>Nyctalus lasiopterus</i>						++++
<i>Nyctalus leisleri</i>			+			+++
<i>Nyctalus noctula</i>	+	+	++	+	+	++++
<i>Otonycteris hemprichii</i>						
<i>Pipistrellus kuhlii</i>	+	+	++++	+	+	+
<i>Pipistrellus nathusii</i>		+	+++	+	+	+++
<i>Pipistrellus pipistrellus</i>	+	++	+++	+	+	++
<i>Pipistrellus pygmaeus</i>	+	+	++	+	+	++
<i>Plecotus auritus</i>	++	+++	++	+		++
<i>Plecotus austriacus</i>	++	+++	++	+		+
<i>Plecotus kolombatovici</i>		+++	+++			
<i>Plecotus macrobullaris</i>		++	++	+		
<i>Plecotus sardus</i>			+++			
<i>Rhinolophus blasii</i>						
<i>Rhinolophus euryale</i>	+	+				
<i>Rhinolophus ferrumequinum</i>	++	++	+	+	+	
<i>Rhinolophus hipposideros</i>	++	++	++	+	+	
<i>Rhinolophus mehelyi</i>						
<i>Rousettus aegyptiacus</i>						
<i>Tadarida teniotis</i>			+		+	+
<i>Taphozous nudiventris</i>						
<i>Vespertilio murinus</i>	+	+	+++	+		+

High dependence in: 1 – 20% of countries (+); 21 – 40% (++); 41 – 60 % (+++); 61 – 80 % (++++); 81 – 100 % (+++++). Countries that could not specify a degree of dependence (answers “not known” or “bat species present”) are excluded.




Figure 1. Percentage of European bat species dependent on different overground roost types (species with high or medium dependence in at least one country were included; see also Annex 2).

3.2 Geographic pattern of dependence of bat species on overground roost types

The questionnaire data merits further analysis which is largely beyond the scope of this report. Nonetheless, some interesting patterns are obvious from the preliminary analyses presented in the figures below. Not surprisingly, it is apparent that in northern European countries a high percentage of the bat fauna rely on roosts in buildings such as castles, churches, houses and barns, compared to the percentage seen in the southern countries (see Figure 2). It would also appear that churches and houses are important for bats throughout Europe (Figures 4 and 5 respectively), whereas barns and bridges are only used in certain countries (Figures 6 and 7 respectively). To some extent, of course,

this is a reflection of the research that has been carried out; as mentioned previously, tree roosting species have not been widely studied and consequently the importance of trees is probably underestimated for many countries (Figure 8).

By and large, a comparison of bat dependence on specific roost types in neighbouring countries provides a coherent picture (e.g. Figure 4). When compiling data, best expert judgement should always be used. Where big differences between neighbouring countries do occur (e.g. Figure 8), they can often be explained by the lack of reliable data on a country level, leading national experts to adopt a cautious approach and declare dependence of particular bat species on particular overground roost types as “unknown.”




Figure 2. Percentage of bat species highly dependent on overground roosts in potential cultural heritage buildings (castles, churches, houses and barns combined). (Only bat species mentioned by MITCHELL-JONES et al. (1999) are considered.)

Figure 2 shows the percentage of bats across Europe which is highly dependent on overground roosts in potential cultural heritage buildings (castles, churches, houses and barns combined). Figures 3-8 present the absolute number and percentage of bat species highly dependent on specific overground roost types in each country.

3.3 Intraspecific variation across the European range

It is clear from the responses to the questionnaire that while certain bat species can be found in the same type of overground roosts across their range, other bats show marked variation in their roost choices across Europe. To some extent this reflects the availability of specific roost types.

Rhinolophus hipposideros provides a good example of this. Churches are highly important for this species in Austria, Slovenia and Slovakia and are of medium importance in neighbouring Hungary, Czech Republic, Germany and France. Further south and east, in Serbia, Montenegro, Bulgaria, Ukraine, Russia and Georgia, churches are less important for this species. Much of this variation can probably be attributed to differences in church construction. In general, catholic/evangelic churches, which predominate in western and central Europe, have large accessible attics suitable for use by *R. hipposideros*. Large attics are not so common in the orthodox churches further east. In contrast to this general trend, the churches in Ireland and in parts of the Mediterranean tend not to have attic spaces,

and when they are present they usually do not have openings large enough for *R. hipposideros* to use. Consequently, in these areas, this species is seldom found roosting in churches, but uses houses and barns instead. In general, houses and barns are often very important for *R. hipposideros* where churches and castles are not. Figure 9 illustrates this further.

Another good example of this changing dependence on overground roosts types

is *Myotis myotis*. In Bulgaria, Romania and Serbia nursery roosts for *M. myotis* are predominantly in caves. In the southern parts of Slovenia, nursery groups can still be found in caves, however, in northern Slovenia and further north again, in Austria and Germany, all nursery roosts of *M. myotis* are located in buildings (Figure 10). Similar clinal (south to north) changes of *M. myotis*' dependence on overground roosts can be expected in other parts of Europe as well.




Figure 4. Absolute number and percentage of bat species highly dependent on roosts in churches in EUROBATS range states.




Figure 3. Absolute number and percentage of bat species highly dependent on roosts in castles/fortifications in EUROBATS range states.




Figure 5. Absolute number and percentage of bat species highly dependent on roosts in houses/blocks of flats in EUROBATS range states.




Figure 10. Estimated dependence of *Myotis myotis* on roosts in potential cultural heritage buildings (castles, churches, houses and barns combined).

4 Roost protection

Three main forms of protection for overground roosts can be recognised: legal protection, physical protection and education/information.

4.1 Legal protection

Most EUROBATS range states have some form of national legislation protecting bat roosts, although a small number do not. Furthermore, specific legislation applies to the 27 EU Member States – in particular all microchiroptera species are listed on Annex IV of the EU Habitats Directive (92/43/EEC). The full text of this Directive can be found at: http://ec.europa.eu/environment/nature/legislation/habitatsdirective/index_en.htm.

Article 12(1) of this Directive requires Member States to implement a system of strict protection. 12(1) b) and 12(1) d) are particularly relevant, they prohibit:

“b) deliberate disturbance of these species, particularly during the period of breeding, rearing, hibernation and migration; (...)

d) deterioration or destruction of breeding sites or resting places.”

It is worth noting that the transposition of this Directive into national law can lead to some variation in implementation between countries. However, the European Commission has produced a guidance document to help clarify the actual interpretation of Article 12. This document includes commentary on many issues of direct relevance to bat protection, including definitions of resting places, guidance on when roost deterioration/destruction may be exempt from the provisions of the Directive *etc.* It is recom-

mended reading and can be downloaded for free from the Commission's website: http://ec.europa.eu/environment/nature/conservation/species/guidance/index_en.htm.

The Convention on the Conservation of Migratory Species of Wild Animals (CMS, also known as the Bonn Convention) was instigated in 1979 in recognition of the fact that migratory animals can only be properly protected if conservation activities are carried out over the entire migratory range of the species. The EUROBATS Agreement was set up in 1994 under this Convention. This Agreement aims to protect all species of bats identified in Europe through legislation, education, conservation measures and international co-operation between Parties and with Non-Party Range States.

Across Europe, bats are further protected under the Convention on the Conservation of European Wildlife and Natural Habitats (Bern Convention 1979). All bat species are listed on Appendix II (Strictly protected fauna species), except *Pipistrellus pipistrellus*, which is listed on Appendix III (Protected fauna species).

Bat roosts can also be afforded effective legal protection through NGO or State ownership. In these cases particular buildings may be bought or leased for the specific protection of an important bat roost, or with roost protection being one of a number of conservation goals. For example, a registered charity called The Vincent Wildlife Trust manages over 50 reserves for horseshoe bats across England, Wales and Ireland.

4.2 Physical protection

Overground roosts can be protected from disturbance by a number of means, including grilling, fencing and blocking off. All in effect restrict public access. The statutory nature conservation authority should always be consulted before such measures are applied. In all cases it is important that these protective measures should not have any accidental adverse impact on the bats themselves; the frequency of comings and goings and the number of bats using an individual roost should always be considered. It should also be remembered that different species could respond in different ways to certain measures, *e.g.* grilling of cave entrances can have a negative impact on *Miniopterus schreibersii* populations, so fencing may be more appropriate for this species.

Wherever possible, the necessary works should be timed to avoid disturbing the bats. Extensive details on physical protection methods can be found in MITCHELL-JONES *et al.* (2007); many of these approaches are equally relevant to overground roosts. Numerous examples of physical protection measures for bat roosts in buildings are also assembled in MITCHELL-JONES (2004), MITCHELL-JONES & McLEISH (2004), THE NATIONAL TRUST (2001), SARGENT (1995), SIMON *et al.* (2004) and in REITER & ZAHN (2006).

4.3 Education / information

The availability of readily accessible and practical information is key to the protection of bat roosts. This can include web-based resources as well as published materials and telephone helplines. While generic guidance is valuable, for maximum effect, focussed information should also be avail-

able, *e.g.* information on bat friendly bridge repairs for local authorities and information on bats in churches for church authorities.

4.3.1 Websites

Web-based information sources are becoming more common. They allow easy and free access to the latest information on best building practice as well as bat ecology and biology from a wide range of countries and in different languages. The availability of this information can be particularly useful for conservationists with limited personal experience of conservation measures for bats in different kinds of buildings. It should always be remembered, however, that situations will vary between bat roosts and even the same species can have different requirements in distant parts of its range.

Non-governmental organisations (NGOs) and academic research groups play an important role in the area of bat education and the provision of related information throughout Europe. They often act as a focal point for frequently asked questions from the public and media. See for example the websites of:

- The Dutch Mammal Society: www.vzz.nl;
- The Italian Chiroptera Research Group: <http://biocenosi.dipbsf.uninubria.it/chiroptera/>;
- The Russian Bat Research Group: <http://zmmu.msu.ru/bats>;
- Bat Conservation Ireland: www.batconservationireland.org;
- SFEPM in France: <http://www.sfepm.org/chiropteres.htm>.

The EUROBATS website provides an extensive list of links to bat conservation organisations across Europe (http://www.eurobats.org/links/links_country.htm).

A recent INTERREG III B project in Austria and Germany provides a good example of the value of publishing project results on the web. The Co-ordination Centre for Bat Conservation and Research in Austria (KFFÖ) and the Co-ordination Centre for Bat Conservation in South Bavaria looked in detail at the effects of building renovation works on various bat species. The resulting report, *Guidelines for the renovation of buildings hosting bat roosts in the Alpine area/Leitfaden zur Sanierung von Fledermausquartieren im Alpenraum* (REITER & ZAHN 2006), includes, on a species by species basis, an examination of critical factors that have to be considered before, during and after renovation works. The report can be downloaded in German or in English from www.fledermausschutz.at/Sets/Literatur-Set.htm (section "download").

Some other very useful general publications available on the web include:

- MITCHELL-JONES (2004): *Bat mitigation guidelines* (<http://naturalengland.etraderstores.com/NaturalEnglandShop/Product.aspx?ProductID=77002188-97f9-45a5-86a6-326a7ea3cd69>);
- MITCHELL-JONES & McLEISH (2004): *The bat workers manual* (www.jncc.gov.uk/page-2861);
- KELLEHER & MARNELL (2006): *Bat mitigation guidelines for Ireland* (<http://www.npws.ie/en/media/Media,4981,en.pdf>).

4.3.2 Telephone helplines

The personal approach in providing advice is often vital, particularly in emergency situations where immediate action is threatened or required. Helplines are available



Figure 11. Example of bat roost notice in situ, Dublin, Ireland.
© F. Marnell

in many countries, often run by NGOs and sometimes through direct or indirect Government funding. If you don't have a helpline in your country, you may find helplines in adjacent countries that can offer general advice.

4.3.3 Site notices

Although conservationists in general often prefer to keep the location of important, unprotected sites secret, sensible use of site notices can be an effective way of alerting developers to the importance of a building or bridge for bats. The notice might usefully state the legal provisions under which bats are protected and provide contact details for the relevant statutory nature conservation authority or NGO as appropriate. If the site is only important for bats for part of the year, the notice could explain this as well.

5 Challenges posed by roosts in domestic dwellings



Many species of bats have adapted to living, for at least some part of the year, in houses, flats or other forms of residential buildings. In many cases, their presence can go undetected by the residents and the roost may only be discovered when renovation work is proposed. The ready availability of practical advice can offset many potential problems and allow the homeowner to plan the works in such a way as to avoid impacting the bats.

On occasion bats roosting in a domestic dwelling can cause a nuisance and some form of intervention may be needed. Once again, the ready availability to homeowners of free, practical advice at this point is critical. In many countries free advice is provided by the statutory nature conservation organisation (SNCO) and/or the local bat group. In some countries such advice is available online in the form of web pages or leaflets and manuals to download. An example of an advice sheet which an SNCO might make available is shown in the box in chapter 5.1.

The best results are often achieved when an informed bat worker is available to visit concerned householders and discuss the perceived problems with them directly. Where this occurs a solution can nearly always be found which satisfies the homeowner and secures the bats' future. This approach is well established in some countries, for example in the UK

where the SNCOs and voluntary bat workers together provide a free support system for householders with bat concerns. Table 2 provides examples of some of the most

common situations that arise when bats are discovered in houses and suggests possible solutions.

Table 2. Bats in houses: common scenarios and possible solutions (adapted from Natural England).

Scenario	Possible solutions
The householder complains of a serious smell of bats or the noise from the roost has kept family members awake.	Where a large build-up of droppings has occurred these will need to be removed. Improving ventilation may help reduce a smell problem, though this may need to be followed up by building work. Noise problems can be intractable, but information about the seasonal nature of the disturbance is often sufficient reassurance.
The householder is phobic, expresses a fear of bats or is clearly worried about the continuing presence of the bats. This could include a strong aversion to accumulated bat droppings on the outside of the property, though there is no evidence that these present a real disease threat.	Reassurance by an expert is helpful for many people, though it needs fine judgement to be certain the householder has changed his views. Droppings problems may be solved with deflector boards.
Accumulated bat droppings and urine are causing damage to the building, most often by causing internal staining. This problem may be coupled with a smell problem.	This is relatively rare in domestic dwellings. Most householders can be persuaded to retain the bat roost if the droppings can be removed and remedial works can prevent a recurrence of the problem. Installing impermeable barriers can sometimes work.
Bats are found by the householder in the living area of the house. The most common situation is baby or inexperienced bats crawling from their roost area into the living area. If these bats are not rescued from within the living area they are likely to die of dehydration or starvation.	The primary action is to locate and block the internal point(s) of entry of the bats from their roost area to the living area of the house to prevent a recurrence of the problem. In some cases, perhaps because of the age or construction of the building, it may be more effective to limit the roosting areas of the bats. This could involve creating an internal bat house or blocking off parts of the attic/cellar space. The SNCO should be contacted for advice in these cases.
The householder dislikes bats, but is not afraid of them and the bats are not causing any noise, smell or damage problems.	Reassurance including information about bat biology (e.g. seasonal use of roost; only one young per year) is often all that is required.
The householder wishes to sell the house without a bat roost.	
The householder wishes to undertake remedial timber treatment in a private house for maintenance, repair or refurbishment purposes.	Carry out the work at a time of year when bats are least likely to be present. Use a product that is not known to be harmful to bats.
The householder wishes to undertake repairs or redecoration of gutters, soffits etc. close to the entrance to a bat roost.	Carry out the work at a time of year when bats are least likely to be present. Ensure that roost entrances are not blocked or altered.
The householder wishes to undertake reroofing for repair or refurbishment.	Carry out the entire work programme at a time of year when bats are least likely to be present. Ensure that roost entrances are not blocked or altered and that the places where bats roost are not altered. Ensure that any roof coverings, including underfelt, are appropriate.
Treatment of cluster-fly infestations.	Carry out any insecticide treatment at a time of year when bats are least likely to be present. Use a product not known to be harmful to bats.

Figure 12. Block of flats in Slovenia where a roost of *Nyctalus noctula* under the exterior cladding only became apparent when renovation work commenced. © P. Presetnik



5.1 Disturbance or exclusion

In very limited circumstances there may be a genuine reason why a bat roost cannot be accommodated at its current location. In EU countries, where all bat species and their roosting places are protected under the Habitats Directive, disturbance to bat roosts or bat exclusions can only be carried out under licence. These licences (or derogations, as they are referred to under the Directive) must be processed by the SNCO and can only be granted subject to the parameters outlined in Article 16 of the Habitats Directive. As the Directive is implemented in each EU country through national regulations there may be some variation in interpretation. However, in general, such licences can only be issued provided that:

- there is no satisfactory alternative **and**
- there is no impact on the conservation status of the species **and**
- the work is for imperative reasons of overriding public interest **or** has conse-

quences of primary importance for the environment **or** serves a public health and safety purpose.

Non-EU countries may have similar legislation and householders should always contact their SNCO before carrying out activities that may impact on bat roosts. General advice for householders with bats exists in many EUROBATS range states and can also be found online in several languages, *e.g.*

- in French at http://www.gmb.asso.fr/les_chauves_souris.html;
- in German at http://www.nabu.de/m05/m05_02/01506.html;
- in Russian at http://zmmu.msu.ru/bats/popular/v_dome.htm.

The EUROBATS website provides a more complete list of bat conservation websites by country at http://www.eurobats.org/links/links_country.htm.

The following English example shows how short and coherent guidance for house owners could be formulated.

Example of guidance for householders with bats

There are 17 different species of bats in England, some very rare, others still quite widespread. These fascinating mammals are heavily dependent on buildings as they often use them as roosts at different times of the year. Bats rarely cause any problems when they roost in houses, but if you are worried about their presence or you want to do something that would affect them or their roosts you should contact Natural England for advice. Many householders have lived happily with their bats for many years. More information about bats and their conservation is given in our free booklet *Focus on Bats* or you can contact Natural England or the Bat Conservation Trust for advice.

Bats and the law

Because populations of most species have declined in past decades, all British bats have been protected by law since 1982. The legal protection they receive has recently been strengthened by changes to the law arising from European Union obligations.

In summary, you may commit a criminal offence if you:

- Deliberately capture, injure or kill a bat;
- Intentionally or recklessly disturb a bat in its roost or deliberately disturb a group of bats;
- Damage or destroy the breeding or resting place (roost) of a bat;
- Possess a bat (alive or dead) or any part of a bat;
- Intentionally or recklessly obstruct access to a bat roost.

Working within the law

There are two main ways in which householders who have bats roosting somewhere within their property can work within the law:

1. Avoid committing offences

It is always preferable to avoid disturbing bats or damaging their roosts if at all possible. That way no offence is likely to be committed and you help with the conservation of these threatened species.

Repairs, maintenance or refurbishment

If you want to carry out repairs, maintenance or refurbishment of your house and believe this might affect the bats or their roost you are advised to consider how you can modify the way you carry out the work so as to avoid committing an offence. This may involve carrying out the work at a particular time of year or using particular materials or methods. You will need to take care that the access points used by the bats are not blocked and that their roosting areas are not damaged as this would be an offence even if it was not intentional. You are strongly recommended to contact your local Natural England office for free advice before you begin the work. For minor works or maintenance, we will usually suggest that a local volunteer or member of staff visits you in order inspect the situation and advise on how best to proceed without breaking the law. We will then confirm this advice in writing. If the work cannot be carried out without affecting the bats or their roost, you are likely to need a licence, as described below. There is, however, no guarantee that a licence will be granted.



Concerns about bats

If you find bats roosting in your property and are concerned about them, please contact your local Natural England office or the Bat Conservation Trust for free advice. We will usually suggest that a local volunteer or member of staff visits you in order to assess the situation and advise on how any problems can be resolved. If necessary, our representative will explain what remedies are available and assist you in deciding on a suitable course of action. We strongly recommend that you do not take any action until you have received advice as this could result in an offence being committed.

2. Work under a licence

In some circumstances, Natural England can issue licences under the Habitats Regulations to permit what would otherwise be illegal actions. However, the reasons for which we can issue a licence are limited in law and, furthermore, we can only issue one if you can show there is no alternative way of carrying out the work and that the work will not adversely affect the local bat population. It is generally far better to try to avoid affecting the bats or their roost wherever possible, as described above. If you believe or have been informed that a licence will be needed, further information about the application process is available on our website or from our local office.

One situation in which a licence is likely to be needed is where repairs, refurbishment or development within or adjacent to your property, such as a loft conversion or extension, will unavoidably damage or destroy bat roosts. In these circumstances, you should still seek advice from Natural England, but we are likely to recommend that you employ a professional ecological consultant to assist you with the licensing process.

Contacts and further advice

Natural England: www.naturalengland.org.uk

Bat Conservation Trust: www.bats.org.uk

Text adapted from Natural England

6 Focus on buildings of cultural heritage

As a general principle, older structures can support a greater variety of bat species than newer ones. Thus, buildings of cultural heritage importance such as castles and churches can play a key role in providing roost sites for many of Europe's bat species. In certain areas, bats may occupy the vast majority of older buildings. In one German study, evidence of use by bats was found in 80% (46/55) of the church attics investigated (SIMON *et al.* 2004).

General features of older buildings that make them attractive to bats include the greater use of natural stone and large hardwood timbers, a wide range of constructional features, limited human disturbance and a certain amount of weathering (HUTSON 1995). It is also a notable feature that bats show a greater degree of site fidelity in old buildings than they do in modern structures, although to some extent this has as much to do with the species concerned as the buildings themselves.

Land use around a building can help determine whether or not the structure itself will be suitable for bats. Frequently, the landscape around heritage buildings is managed in a traditional way that helps to retain features suitable for bats and their insect food (such as trees, permanent pasture and water bodies).

When conflict arises between the protection of bats and of buildings of cultural heritage importance it is usually in one of two ways: either restoration/renovation works are planned for the building that will

impact on the bats, or the bats themselves are causing a disturbance or damage within the building. In some cases these conflicts may be supported by opposing legislation, with the bats being protected under wildlife law and the building and/or its contents protected under other heritage legislation. However, there are many examples to show that such conflicts can normally be resolved to the satisfaction of both the built heritage and the natural heritage.

6.1 Protection of cultural heritage

There are numerous international agreements, treaties and conventions devoted to the protection of our cultural heritage. One of the leading organisations in this area is the International Centre for the Study of the Preservation and Restoration of Cultural Property (ICCROM) in Rome, which is an intergovernmental organisation with a membership of over 100 countries. The decision to found ICCROM was made at the 9th UNESCO General Conference in New Delhi in 1956, at a time of mounting interest in the protection and preservation of cultural heritage. The Centre was subsequently established in Rome in 1959 at the invitation of the Government of Italy. It is the only institution of its kind with a worldwide mandate to promote the conservation of all types of cultural heritage, both movable and immovable. ICCROM aims at improving the quality of conservation practice as well as raising awareness about the importance of preserving cultural heritage.

UNESCO's Convention Concerning the Protection of the World Cultural and Natural Heritage, signed in Paris in 1972, recognised the dual need for protection of both natural and built heritage elements. Further information on this agreement can be found at http://portal.unesco.org/culture/en/ev.php-URL_ID=8453&URL_DO=DO_TOPIC&URL_SECTION=201.html.

In many countries, the responsibility for the protection of natural heritage and the conservation of the built heritage fall under different government departments. The need to address the issues relating to bat conservation in historic buildings through cooperation between separate government sections has been recognised recently in Italy. In 2006, a joint project was initiated there by the Ministry of Environment and the Ministry of Culture. This project will lead to the creation of a database of bat roosts in heritage buildings as well as the preparation of guidelines for public works in historic buildings. (A report on this project can be downloaded at <http://fauna.dipbsf.uninsubria.it/chiroptera/buildings/index.html>.)

Similarly in 2006, a harmonised database of bat roosts and buildings of cultural heritage was commissioned by the Slovenian Ministry of the Environment and Spatial Planning.

There are many examples from throughout Europe to show how bats need not be impacted during building works (see case studies below). Indeed, with some careful planning, the status of bats in a building of cultural heritage can often be enhanced during such operations. Equally, it has been shown that if bat expertise is involved from the early planning stages of a restoration project, and a flexible approach is taken to

the scheduling of the works, the bats can be satisfactorily accommodated throughout the project at little or no additional cost and without compromising the aims of the works.

6.2 Damage by bats in buildings of cultural heritage

Bats flying around within an occupied building can sometimes be a cause of disturbance or concern. Furthermore, bat excreta may cause damage to vulnerable objects and furnishings in buildings.

Droppings, over a protracted period of time, may cause pitting, long-term staining and etching to porous materials such as painted wall surfaces, wooden monuments and stone sculptures. Bat urine (which is 70% urea) is chemically more aggressive and therefore of even greater conservation concern. It can cause spotting and etching of wooden, metal and painted surfaces (PAINE 1993 & undated).



Figure 13. Damage to stonework at Penmon Priory, Wales, United Kingdom, due to long-term exposure to bat urine.

© J. Matthews

Before any management of these situations begins it is essential to assess bat activity and its effects on the building's contents. In most cases, there are practical steps that can be taken to manage these problems without compromising the status of the bats or the cultural heritage.

In each individual case, information will need to be gathered on the bats themselves; the rate of deposition and the seasons when it occurs; the area/articles being damaged; and the extent of the damage. Once these assessments have been carried out an informed decision can be made on which of the following management techniques may be most usefully implemented (these are adapted from advice prepared by S. PAINE (undated) for English Heritage):

- **Do nothing** – Bats are usually only present in buildings for a limited season. Normally summer colonies consist of females which have gathered to give birth and rear their single young. They may not be a problem if they occur in very small numbers or only use parts of a building without vulnerable or significant objects.
- **Remove droppings** – Large accumulations of droppings (e.g. in attics) can give rise to concerns about smell, health and safety. However, removing the droppings is normally quite straightforward, providing there is access to the roof space, and the droppings make good garden fertiliser (see case study 1 below and another case study in MITCHELL-JONES & McLEISH 2004, p. 85).
- **Moving objects** – If an object being exposed to bat excreta is freestanding, it may be possible to move it to a location with a lower rate of deposition.
- **Covers** – Covers may be appropriate when deposition is localised or if there are

few vulnerable objects. They are not suitable if deposition occurs throughout a room, as there would be a great aesthetic impact. Porous materials such as linen or natural carpet are suitable covers; however, polythene should not be used as this may create a moist microclimate around the object. Covers need only be used during the period when bats are present and can be removed for exhibitions, services etc.

- **Coatings** – Synthetic lacquers offer some protection against bat damage and may be acceptable on historically and artistically insignificant metal and wooden objects. Natural organic coatings (such as beeswax) offer little protection against bat urine.
- **Deflector boards** – A wooden board 100-150 mm wide and 1-2 m long can be positioned at an angle beneath a roost or access point to deflect and/or catch any droppings. This can be useful to reduce rates of deposition in specific areas. The board can be erected for the summer and removed at other times of the year for cleaning.
- **Relocation of roosts or access points** – This has been used with some success in the past. Excluding bats from one roost site will reduce the impact in the immediate area, but may cause them to move to another part of the building and have an undesirable effect there. This can be avoided by blocking off potential roost sites first. Relocation should be considered carefully, with the relevant Statutory Nature Conservation Organisation (SNCO) and bat experts or local bat group being contacted for advice and permission. However, some species, e.g. *Myotis myotis*, can be reluctant to use new sites or even new entrances to the same roost (REITER & ZAHN 2006, BIEDERMANN *et al.* 2008).

• **Exclusion** – This decision, only to be taken by the SNCO, will depend upon a variety of criteria, including the value of the object at risk and the rarity of the bat species. Advice and permission should be sought from both the relevant SNCO and those responsible for the conservation of the historic artefacts. Exclusion may be difficult and expensive. The provision of an alternative roost is usually required.

6.3 Accommodating bats during renovation/restoration

6.3.1 Avoidance (adapted from MITCHELL-JONES 2004)

The most common and effective method of minimising the impact of renovation or restoration works on bats is to carry out

the work at an appropriate time of the year. More than half of respondent countries to the overground roost questionnaire had employed this approach. The great majority of roosts in buildings are used only seasonally, so there is usually some period when bats are not present. Although there are differences between species, maternity sites are generally occupied between May and September and hibernation sites between October and March, depending on the weather and geographical area. An adequate survey and good understanding of the seasonal activity patterns of the particular species involved will help in determining the optimum time to carry out the proposed work. The recommended times shown in Table 3 should be modified in the light of

Case study 1: Ratková Church, Slovakia

The loft of the Lutheran church in the village of Ratková, Slovakia, is occupied by a nursery colony of *Myotis myotis* and *Myotis blythii* in summer. The colony was discovered in 1992 and is the biggest colony of this type known in Slovakia, with up to 5,000 individuals present. A thick layer of bat guano had accumulated below the colony over the years; in places the layer of guano exceeded 1 m. The weight of the guano was about 10 tonnes, giving rise to concerns about the ceiling of the church.

On 3-4 December 2004, the loft of the church was cleaned with the help of the employees of the Muránska Planina National Park and Slovak Bat Conservation Group (SON) members. The guano was

bagged and distributed to members of the local community as fertiliser. The colony continues to thrive and the ceiling of the church is no longer threatened with collapse.

See SON website for further details of this work: <http://www.netopiere.sk/aktuality/2004/12/03/cistenie-kostola-v-ratkovej>.

Figure 14. Ratková Church, Slovak Republic, clean-up team with bagged bat guano.
© Slovak Bat Conservation Group



Table 3. Optimum season for works in different types of roosts (modified from MITCHELL-JONES 2004). The period of works may be extended if the way in which the bats use the site is well understood.

Bat usage of site	Optimum period for carrying out works (some variation between species, and geographical regions)
Maternity	1 October – 1 April
Summer (not a proven maternity site)	1 September – 1 May
Hibernation	1 May – 1 October
Mating / swarming	1 November – 1 August

site-specific species information. For example, some species, most notably *Plecotus auritus* and *Rhinolophus hipposideros*, tend to remain in summer sites until well into autumn or even winter, so care may be needed when drawing up works timetables where these species are present.

Bats are most vulnerable in buildings during the summer, when large numbers may be gathered together and young bats, unable to fly, may be present. Operations to known breeding sites should therefore be timed to avoid the summer months. Very large rebuilding or renovation projects may take many months to complete and may need to continue through the sum-

mer, which is the favoured season for re-roofing. The best solution in such cases is to complete and secure the main roosting area before the bats return to breed. If this is not possible, work should be sufficiently advanced by May or June for returning bats to be dissuaded from breeding in that site for that year. As part of the mitigation, alternative roosts appropriate to the species should be provided in a nearby location. Another possible solution is to divide the roof with a temporary barrier and work on one section at a time. This procedure has been used successfully on a number of occasions (e.g. REITER & ZAHN 2006).

Case study 2: St Cadoc's Church, Wales, United Kingdom

Three species of bat roost in St Cadoc's church in Wales – *Rhinolophus hipposideros*, *Plecotus auritus* and *Myotis nattereri*. The church dates back to the early 1200s and is of considerable historical importance. In 2002 it was discovered that essential repairs were required and scaffolding was immediately erected to stop the roof from collapsing. No further works were carried out, however, until the bats had left the church in the autumn. At that stage a polythene tent was constructed over the roof to allow the restoration works to be carried out over the winter months with a view to having the building ready for occupation by bats again the following spring.

The work took longer than expected, however, and the builders were still on site when the bats returned. Through agreement with the local SNCO (here the Countryside Council for Wales), work was able to continue on the main roof, but the tower where the bats roosted was left untouched during the breeding season. The bats successfully reared their young, despite the ongoing restoration works nearby, and the remaining works were completed over the autumn.

The peak numbers of lesser horseshoe bats, which have been counted at the site each year, appear to indicate no negative impact of the building work:

2001: 140;
2002: 133;
2003: 152;
2004: 116;
2005: 126;
2006: 142;
2007: no data;
2008: 180.

Count data from Monmouthshire Bat Group.

For further information please contact the Countryside Council of Wales at Enquiries@ccw.gov.uk.

Figure 15. St Cadoc's Church, Wales, United Kingdom, undergoing essential repairs.
© C. Roberts



Where the same structure is used by bats throughout the year, the optimum time for works of all types is likely to lie outside the main breeding season (to avoid times when females in later stages of pregnancy and non-flying babies may be present) and outside the main hibernation season (to avoid

times when disturbance may impact on survival, or bats may not be sufficiently active to get out of the way). Spring and autumn generally provide the optimum periods for such operations.

In spring and autumn bats will be able to feed on most nights, but may be active or

torpid during the day, depending on weather conditions. Active bats will usually keep out of the way of any operations, but torpid bats may need to be gently moved to a safe place, preferably without causing them to fly out in daylight. Wherever possible, the objective should be to persuade bats to move of their own accord and they should be physically moved only as a last resort.

In many cases it is not easy to determine if a building is used for hibernation, except occasionally in the case of lesser horseshoe and long-eared bats in cellars. Where bats are known to be present, significant disturbance during the winter must be avoided and work should be delayed until after hibernation if possible. Repeated disturbance to bats during the winter can seriously deplete their food reserves.

If there are overriding reasons for carrying out works during a sensitive period, for example in roosts that are used throughout the year, it will be necessary to structure and time the works so as to ensure that the bats always have some undisturbed and secure areas. This may involve the installation of temporary partitions and adopting working practices that minimise disturbance to sensitive areas.

6.3.2 Incorporating existing roosts into renovated buildings (adapted from MITCHELL-JONES 2004)

The renovation of heritage buildings used by bats can provide opportunities to incorporate existing roosts into the final structure. Apart from the timing of the works, the two most critical issues in maintaining a roost *in situ* are the size and suitability of the final roost and the disposition of the entrances and flight paths, including the location of any exterior lighting or vegetation.

6.3.2.1 Roost size

The size of roost required depends on the species, as some require voids sufficiently large to fly into whereas others are more likely to roost in crevices and use direct exterior access. In addition, some species may require light-sampling areas where they can fly in and out before finally emerging. Hibernation roosts in buildings are normally underground. Table 4 gives an indication of summer roost preferences for some species, though there is a great deal of variation; the overall objective should be to maintain the roost size as close to the original as possible.

Case study 3: Grad na Goričkem, Slovenia

Grad na Goričkem lies in northeastern Slovenia, close to Austria and Hungary. It is a castle of cultural heritage importance dating from the middle ages. When plans were developed to transform the castle into a visitor centre for cross-border landscape parks, it provided an opportunity to improve the roosting habitat of the castle's bats.

Bats were first discovered in the castle in 1999. Intensive research followed on the composition of the bat fauna, seasonal dynamics of species and the microclimates of the areas being used by bats. Volunteer involvement was also important in developing an understanding of the importance of the building for bats. Conservation work was then undertaken to protect the bats from disturbance. Funding was provided by the State and also through an INTERREG IIIA project (Conservation of amphibians and bats in the Alpine & Adriatic region).

Ten bat species (one third of all Slovenian species) were found to use the site; the cellars provide hibernation sites for *Rhinolophus hipposideros*, *Myotis myotis*, *Barbastella barbastellus* and even occasionally for *Myotis bechsteinii*. *M. myotis* use the cellars as mating quarters as well. Up to 100 *Miniopterus schreibersii* have been recorded in the castle, making it one of the biggest known roosts for this species in the northwestern part of the Pannonian basin. *R. hipposideros* also forms a small nursery group in the attic of the castle. As underground habitats are generally rare in the region, the cellars are thought to be an important swarming site for bats in the wider area.

The building works required the complete demolition and reconstruction of parts of the castle used by bats. On the basis of the research, mitigation measures were recommended during the renovation, includ-

ing the designation of part of the cellars as a bat roost. Extensive discussion took place between nature conservation and cultural heritage officers to agree the position and size of a new entrance for bats (Figure 16). Follow up monitoring is now required to ensure that the conservation measures are effective, but it seems that the conservation efforts to date have been successful.

For further details of this work see KRAINER *et al.* (2007).



Figure 16. Northwest tower of Grad na Goričkem castle, Slovenia, during and at the end of reconstruction in year 2006 (circles mark new entrance openings for bats).

© P. Presetnik

Table 4. Specific roost types and sizes for bat species highly dependent on roosts in buildings (compiled from LIMPENS *et al.* (2000), MITCHELL-JONES (2004), REITER & ZAHN (2006), SIMON *et al.* (2004) and answers from IWG5 questionnaires, EUROBATS national reports and personal unpublished data; see also Table 1).

Species	Summer / maternity roosts
<i>Barbastella barbastellus</i>	Crevice dweller; may require light-sampling areas. Roosts behind window shutters, behind outer wall panelling and similar crevices.
<i>Eptesicus nilssonii</i>	Crevice dwellers; often in houses.
<i>Eptesicus bottae</i>	
<i>Eptesicus serotinus</i>	Roosts in roof spaces. However, the hanging places are mostly well hidden in crevices (chimney breasts, ridge boards <i>etc.</i>) as well as behind wall facades or in roller shutter boxes.
<i>Miniopterus schreibersii</i>	In the northern part of its range, some nursery groups are found in church lofts or towers; requires large entrance openings.
<i>Myotis alcaethoe</i>	Crevice dweller.
<i>Myotis brandtii</i>	Crevice dweller, but may enter roof voids and fly around. In buildings, hanging places are encountered particularly in lofts, although the animals populate the nooks and crannies (e.g. in false ceilings, gaps between beams or between metal sheeting and wall, as well as holes in beams).
<i>Myotis blythii</i>	Nursery roosts are located in larger roof spaces (such as lofts and towers of churches), and more rarely also in bridges and (heated) subterranean spaces. The nursery communities mostly hang free in the roosts and are rarely hidden in crevices
<i>Myotis myotis</i>	
<i>Myotis dasycneme</i>	Summer roosts in lofts and cavity walls of buildings, probably also in hollow trees. May also use bat boxes.
<i>Myotis daubentonii</i>	Hole dweller; may enter roof voids and roost at apex. Relatively rare in houses sometimes found in wall facades and behind window shutters or the casings for roller shutters, but may use castles, tunnels <i>etc.</i> Nurseries and also colonies of males can be found under bridges.
<i>Myotis emarginatus</i>	Nursery roosts are mainly located in roof spaces (e.g. attics and haylofts). In the roosts, the animals mostly hang free, or more rarely are found in confined spaces, such as in mortise joints. They use comparatively bright and only moderately warm spaces that are well structured, for example by having several different levels.
<i>Myotis mystacinus</i>	Crevice dweller, but may enter roof voids and fly around, often located behind vertical outer wall coverings.
<i>Myotis nattereri</i>	Crevice / hole dweller; may require light-sampling areas. Frequent in crevices in timbers in old barns and stables.
<i>Nyctalus leisleri</i>	Crevice / hole dweller; sometimes in buildings, but unlikely to fly inside.
<i>Nyctalus noctula</i>	Hole dweller; seldom found in houses, but can be found in crevices in higher floors of blocks of flats, sometimes also in church attics and bridges.
<i>Pipistrellus kuhlii</i>	Crevice dweller found in wide range of cracks and small hollows, such as shutters, roller shutter boxes, fissures in walls, in wall facades, false ceilings or crevices in the roof area. Winter roosts are found in buildings. These can be in hollow spaces in walls, and their facades.
<i>Pipistrellus nathusii</i>	Crevice dweller.
<i>Pipistrellus pipistrellus</i>	Usually use crevice roosts, such as exterior wall cladding, roller shutter boxes, false ceilings, flat roof cladding, hollow blocks of unplastered house walls, shutters, hiding places in roofs as well as cracks in walls. Additionally, tree hollows and nest boxes are also used as mating roosts, and individual animals can be found in a great variety of hiding places. Does not normally require light-sampling areas.
<i>Pipistrellus pygmaeus</i>	

Table 4 (cont.)

Species	Summer/maternity roosts
<i>Plecotus auritus</i>	Hole dwellers; readily fly within roof voids, churches, private houses. Often in crevices in the roof frames by day, although sometimes in the open.
<i>Plecotus austriacus</i>	
<i>Plecotus kolombatovici</i>	
<i>Plecotus macrobullaris</i>	
<i>Rhinolophus euryale</i>	Horseshoe bats require large roost areas, with flight access into them, where they hang free, e.g. attics of churches, schools, private houses. Normally require associated sheltered light-sampling areas.
<i>Rhinolophus ferrumequinum</i>	
<i>Rhinolophus hipposideros</i>	
<i>Vespertilio murinus</i>	Crevise dweller; usually associated with blocks of flats and private houses.

For species that need to fly within roof voids, notably *Rhinolophus* and *Plecotus* species, it is essential that a sufficiently large space, unobstructed by constructional timbers, is available for the bats to fly in. These species are generally found in older roofs of traditional construction which give a large uncluttered void. Trussed rafter constructions do not provide these conditions. Suitable construction methods are purlin and rafter ('cut and pitch') with ceiling ties or any other construction techniques which are designed to give a large, open roof void. Based on a sample of known roosts, it is unlikely that a void height (floor to ridge board) of less than 2 m will provide sufficient volume, or that an apex length or width of less than 4 m will provide sufficient area. An ideal roof void would have an apex height in excess of 2.8 m and a length and width of 5 m or more (MITCHELL-JONES 2004).

Some recent UK studies on *Myotis nattereri* in barns due for conversion have illustrated some of the difficulties of maintaining appropriate roosts. In these cases, bats were roosting in mortise joints, which

presumably mimic tree cavities, and using the void of the barn as a light-sampling area. In several cases, the bats abandoned the site after conversion, probably because insufficient 'indoor' flight opportunities remained. Full details and recommendations can be found in BRIGGS (2002).

6.3.2.2 Roost entrances

Rhinolophus species and *Miniopterus schreibersii* generally prefer entrances they can fly through (see MITCHELL-JONES & McLEISH (2004), chapter 11 for details and designs), but other species will generally use smaller holes or slits to crawl through. Wherever possible, it is preferable to maintain entrances in their original position so the bats will have no difficulty finding them. Retention of vegetation close to roost entrances can also be important. This provides continuity of flight routes and cover for the bats which protects them from avian predators. External lighting, such as security lights or road or path lighting, close to roost entrances should be avoided (see DOWNS *et al.* 2003, REITER & ZAHN 2006).



Figure 17. Dedicated bat roost entrance, Ireland (also used by lesser horseshoes!). © C. Kelleher

6.3.3 Incorporating new roosts into buildings

The extent to which new roosts can easily be incorporated into new or refurbished buildings depends on the species of bat and the type of building. For those species that require a large roof void to fly in, principally *Rhinolophus* and *Plecotus* species, careful attention must be paid to the design in order to provide a suitable roof void. See above for guidance on roost size and construction and note that trussed rafter construction should normally be avoided. For species that typically roost in crevices, roosting opportunities can be provided in a variety of ways including:

- Access to soffit boxes and eaves via a small gap (15-20 mm) between soffits and wall;
- Timber cladding mounted on 20-30 mm counter battens with bat access at the bottom or sides;

- Access to roof voids via bat bricks, gaps in masonry, soffit gaps, raised lead flashing or purpose-built bat entrances;
- Access to roof voids over the top of a cavity wall by appropriately constructed gaps.

As well as suitable access points, bats also need suitable roosting sites and an appropriate temperature regime.

Most species of bats appear to prefer roosting on timber rather than brick, stone or other similar materials, so the provision of rough timber surfaces may be helpful. Bats may also roost by clinging on to roof lining materials, especially around the roof apex and 1 m or more down the slope. Some types of modern plastic roof linings are too smooth for bats to cling to and should be avoided where possible. If their use is essential, rough timber planks should be placed along the ridge beam to provide roosting opportunities.

For maternity roosts, bats appear to prefer maximum daytime temperatures of between 30° and 40° C, so it is important that the roof receives full sunlight for a large part of the day. This can be assisted if the roof has two ridges at right angles, oriented to capture sunlight throughout the day. As an alternative, a combination of baffles and electric heaters can be used to produce pockets of warm air at the apex of the roof. This technique has been used successfully with *Rhinolophus* bats (SCHOFIELD 2008) and would probably be suitable for other species as well.

Where space permits, large 'bat-boxes' can be built into existing roofs. This approach has the advantage of providing some segregation between the bats and the human occupants of the building. Detailed

guidance is given in the Scottish Natural Heritage (SNH) publication *The design and construction of bat boxes in houses* (SIMPSON & BROWN ARCHITECTS 1996). SNH have recently published a follow up report which includes suggested modifications to previous designs (BAT CONSERVATION TRUST 2006). SIMON *et al.* (2004) provide detailed informa-

tion on the construction of artificial roosts within buildings. They also provide an assessment of the success of these various artificial roost types based on a research programme in Germany. For other practical examples of mitigation measures and alternative roosts see REITER & ZAHN (2006).

Case study 4: Glaninchiquin, Ireland

When an old cottage which contained a nursery colony of *R. hipposideros* was being renovated in south-west Ireland it was not feasible to include a suitable roost in the final design. It was decided to undertake works to an adjacent stone outbuilding to accommodate the bats.

The outbuilding, which was 12 m x 5 m and approximately 10 m from the original cottage, was roofed with slate and had an underlay of mineral felt. A loft was created in the building, with two trap doors and an access point in one of the gables directly into the loft (the original roost had also had a direct gable entrance into a loft). The floor of the loft was insulated to help minimise disturbance as the owner planned to store materials on the ground floor of the outbuilding.

The original roost had a count of 150+ bats in 2003. 130 bats remained in the old roost despite considerable disturbance during 2004. The new roost was constructed at the end of 2004 and by May 2005, the bats had moved in. The peak count for the new roost in 2005 was c. 120. Bats were not able to enter the original roost by this time as all access points had been sealed. The peak count in the new roost in 2007 had risen to 150 animals.

For further information on this case please contact natureconservation@environ.ie.



Figure 18. Outbuilding at Glaninchiquin, Ireland, before and after renovation for *Rhinolophus hipposideros*.

© C. Heardman

One problem with providing roosts in buildings intended as dwellings may be their acceptability to the future inhabitants and for this reason planners and developers are often reluctant to adopt this solution. There is much to be said for providing a dedicated

bat roost as these problems of acceptability can be greatly reduced. MITCHELL-JONES (2004) and SCHOFIELD (2008) provide extensive advice on the design and construction of such dedicated roosts.

Case study 5: Morcegário, Portugal

In 2000, bats were discovered during the environmental impact study for the destruction of a 15-storey building in Portugal. Up to 100 *Tadarida teniotis* and some *Eptesicus serotinus* and *Pipistrellus pygmaeus* were hiding in crevices below concrete plates covering the walls.

Detailed monitoring showed that bats were present in all seasons and favoured walls with higher sun exposure. Bats were present at various heights, but were most abundant above 21 m, where temperatures were warmest. 75% of the bats were found inside crevices less than 3 cm wide.

The developer built a new roost in 2003, 150 m from the original. It was designed, in consultation with the statutory nature conservation organisation, to replicate the original building, although it is only 12 m high. In order to ensure that the thermal characteristics of the crevices were replicated the concrete plates of the original building were re-used. Follow-up monitoring confirmed that the thermal behaviour of the new roost was quite similar to the original one.

To encourage colonization of the new roost, 50 bats were captured and released there when it was finished. The old building was knocked down in 2005. In 2006, 22 *Tadarida teniotis*, 12 *Eptesicus serotinus*

and 4 *Pipistrellus pygmaeus* were recorded in the new roost. In 2007, the maximum numbers seen were 11 *Tadarida teniotis*, 11 *Eptesicus serotinus* and 7 *Pipistrellus pygmaeus*. Monitoring of the new roost is continuing.

For further information on this case please contact site@icnb.pt or rodrigues@icnb.pt.



Figure 19. Old and new *Tadarida* roosts, Portugal.

a) Original building.

b) New roost.

© M. Carapuço

© J. Palmeirim

6.3.4 Barns

Old barns play an important role as roosts for some bat species in certain countries and provide their own challenges when it comes to accommodating bats during renovation or restoration works. A study in the UK has shown that many old timber-framed barns, some dating back several centuries, are now being converted into dwellings. BRIGGS (2002, 2004) found that the vast majority (77%) of converted barns have not maintained their bat species and she questions whether barns with bats should ever be converted. She looked at how bats could

best be accommodated in these conversions and provides details of mitigation measures that should be built into future barn conversion designs. The features covered include:

- Species specific design;
- Roost site retention;
- Light pollution;
- Access;
- Conservation and enhancement of adjacent habitats;
- Timing of the works.

The reader is referred to BRIGGS (2002, 2004) for further details.

Case study 6: Paston Barn, England, United Kingdom

Paston Barn was built in 1581 and is home to a breeding colony of *Barbastella barbastellus* as well as *Pipistrellus pipistrellus* and *Myotis nattereri*. The building is owned by the North Norfolk Historic Buildings Trust, which had initially planned to turn the barn into a visitor centre for the nearby gas works before the bats were discovered. Natural England has since taken a 50-year lease on the building.

The barn has been subject to massive renovation over the last few years, including complete re-thatching, re-pointing, replacement of doors, and restoration of its associated buildings. A steering group, including BCT, Natural England, the local bat group and the Buildings Trust, has controlled restoration work at the site. Measures to minimise the disturbance to the bats have included timing of works outside of the breeding season (though due to delays work sometimes overran), replacing doors with temporary

structures while work was done off site, use of traditional materials and carrying out work by hand where possible. The roost spaces above the lintels, which were favoured by *Barbastella barbastellus*, were maintained and the new doors were constructed to give continued access for the bats.

The collaboration of all parties, and the sensitive nature of the works to date, have ensured the preservation of this historic barn and the continued use of the building by the bats.

Further information is available at <http://www.naturalengland.org.uk/>.



Figure 20. Paston Barn, England, United Kingdom.
© G. Hewson

6.3.5 Bridges

Bridges are not technically buildings. However, bridges are known to be of particular importance for at least 13 species of bats across Europe (see Table 1). For example, out of 328 inspected bridges in Austria 30% were used by bats (PYSARCZUK & REITER 2008). Old bridges, normally made of stone, regularly form part of our cultural heritage. These are subject to different types of disturbance and require different forms of maintenance to other man-made structures which might host bat roosts. Here some general guidelines on the protection of bats in these structures are provided.

In older, smaller stone bridges bats are commonly found in small numbers. A survey of 200 known bridge roosts of *Myotis daubentonii* in Ireland showed that 75% were occupied by 1-5 bats and only 5% held 20 or more bats (SHIEL 1999). Individual bats will use crevices as small as 50 mm deep and 12 mm wide, but larger groups require bigger, deeper roosting sites. Large, concrete motorway bridges with big interiors can provide shelters for many bats (e.g. one of the biggest known maternity roosts of *Rhinolophus hipposideros* in Austria is found in such a bridge).

6.3.5.1 Bridge survey and mitigation measures

Surveys of bridges require a certain degree of expertise. Likely roosts can be identified quite readily, providing there is convenient access to the underside of the bridge, but determining whether they are used by bats is not always easy. The presence of bat droppings may provide a clue, but a fibrescope may be necessary to investigate some cracks. If there is evidence that

a bridge is used by bats then the national nature conservation organisation should be contacted and measures should be taken to ensure that any impact on bats is avoided or, where this is not possible, minimised.

In general the bridge should be as suitable for bats after the required works as it was before. In some cases it may be possible to improve conditions for bats by incorporating specific bat roosts into the structure. As with restoration work of other structures of cultural heritage importance, timing the works to coincide with the period when bats are absent may be sufficient to avoid any impact.

In most cases, the implementation of the following mitigation measures should ensure that bridge renovation works do not negatively impact on bats:

- Careful timing of the works, especially if breeding or hibernating bats roost in the bridge;
- Preserving individual roosting spaces wherever possible;
- Hand pointing in sensitive areas, e.g. around crevices to be retained;
- Creation of new roosts – bat bricks or boxes can be incorporated into a bridge to replace lost crevices.

The Société Française pour l'Etude et la Protection des Mammifères (SFPEM) has produced a useful leaflet (in French) about the use of bridges by bats. The leaflet can be downloaded from <http://www.sfpepm.org/NuitChauveSouris/images2/Savoirplus/plaqponts.pdf>. They estimate that about 10% of bridges in France are used by bats and provide helpful advice on how to accommodate bats in both old and new structures.

Case study 7: Lisconny Bridge, Ireland

Lisconny Bridge is a beautiful five-arch masonry bridge spanning the Unshin River in northwest Ireland. It was built c. 1800 and in 2003 it was determined that the bridge was in need of major strengthening work. There was a large deep fissure running across two of the arch barrels. In addition, the bridge abutments were being seriously undermined by the river and were becoming unstable.

The bridge was known to hold a nursery roost of approximately 25 *Myotis daubentonii*. The roost was located in a deep fissure running across the barrel of one of the arches. Several other crevices were also being used by individual bats. A bat expert was employed by the local authority to work with the local engineer and with the contracting firm. It was agreed that three to four small crevices would be retained under each arch for individual bats. These were ring-marked with white paint before work began. It was also agreed that sections of the large fissure would be left open to accommodate the nursery colony.

Strengthening works commenced at the end of September 2003. This included the laying of new cement floors under the arches, pressure grouting to fill internal voids in the bridge and pressure pointing under the bridge arches. Prior to pressure pointing, all crevices that had been marked for retention were blocked with polystyrene to prevent the infill of grout. Crucially, a fine nozzle was used to apply grout to the undersurface of the arches. The work was done carefully to fill in voids around the stonework so as not to cover the surface of the stones and not to spray over targeted bat crevices. Not only did this allow the bat crevices to be maintained, but it also retained the beauty of the bridge's stonework.

Works were completed by late October 2003. In early July 2004 only four bats were recorded in the nursery crevice. However, most of the other smaller crevices which had been retained held single bats. In July 2005 a nursery colony of approximately 25 bats were recorded in the location where they were first recorded in 1988. Lisconny Bridge shows how bats can be accommodated during bridge strengthening procedures providing there is close liaison between the contractor and the bat specialist both prior to and during strengthening works.

Further information on this case is available at <http://www.batconservationireland.org/pubs/natterer/NattererSummer2005.pdf>.

A further bridge repair case study can be viewed at <http://www.whelan.me.uk/bats/Tattynure/BatsAndDippersInBridges.htm>.



Figure 21. Lisconny Bridge, Ireland, after restoration.
© C. Shiel



Figure 22. Underside of Lisconny Bridge; showing the three sections of the deep fissure that were left open to accommodate the nursery colony of *Myotis daubentonii*. The nursery roost is located in the lowest section. Note how the stonework was carefully grouted around and not over.
© C. Shiel

Similarly DIETZ (2005) provides an overview of the problems and includes well illustrated practical advice in German at <http://www.fledermaus-dietz.de/publications/Dietz%202006%20Fledermaus-Brosch%FCre%20Br%FCcken.pdf>.

6.3.6 Timber treatment, pest control and lead poisoning

Repair and restoration of old buildings often requires timber treatment against infestations of wood-boring insects. *In situ* remedial timber treatment with organochlorine insecticides and some fungicides is thought to be a significant cause of bat mortality across Europe (HERNANDEZ *et al.* 1993, JEFFERIES 1976, RACEY & SWIFT 1986). In recent years, the widespread replacement of certain toxic chemicals, such as lindane, with relatively harmless alternatives (*e.g.* synthetic pyrethroids) has improved the situation for bats. Nonetheless, the guiding principle is that treatment should take place at a time when no bats are present. In most situations, where bats are only present seasonally, this is fairly straightforward. Certain species, however, may be present in buildings all year round and there is no ideal solution in these cases. Advice should be sought from the SNCO.

The control of pest insects or rodents need not lead to any disturbance of bats providing it is done sensitively. Ideally, any treatments would be applied while bats are not using the roost, but localised applications of insecticide powder or rodent poison is unlikely to harm bats. If the control work must be done while the bats are present



Figure 23. Château de Trévez, Brittany, France.
© X. Gremillet (Groupe Mammalogique Breton)

and needs to be more extensive than advice should be sought from the SNCO. Extensive guidance on best practice in the areas of timber treatment and pest control is given in the JNCC's *Bat Worker's Manual* (MITCHELL-JONES & McLEISH 2004).

A particular problem with some older buildings is the existence of lead based paints on girders or other metal structures. Bats can develop lead poisoning by ingesting flakes of this paint during grooming. Such a situation arose in the Château de Trévez in north-west France. The chateau contained a nursery roost of 300 *Rhinolophus ferrumequinum*. Lead and pentachlorophenol poisoning was found to be the cause of high juvenile mortality at the site and in this case it was decided that the best solution was to build a new roost for the bats (GRÉMILLET & BOIREAU 2004, GRÉMILLET 2006).



7 Literature and further reading

- ALLEN, P., I. FORSYTH, P. HALE & S. ROGERS (2000): Bats in Northern Ireland. Their demography as recorded in the historic literature and the data files of the Northern Ireland Bat Group. Special Zoological Supplement to The Irish Naturalists' Journal.
- ALTRINGHAM, J.D. (2003): British bats. Harper Collins, New Naturalist Series, London.
- ANONYMOUS (2003): Natural Heritage and the Law: Bats and people. Scottish Natural Heritage.
- APPLETON, C. (2003): The effect of building works on bats: ten case studies. The National Trust. Available on CD from The National Trust, Conservation Directorate, 33 Sheep St., Cirencester, Glos. GL7 1RQ, UK.
- BALBIERIUS, A. (1981): Šiksnosparniai skrenda per Ventės Raga. Mūsu gamta 9: 16-17.
- BAT CONSERVATION TRUST (2006): A review of the success of bat boxes in houses. Scottish Natural Heritage Report No. 160.
- BIEDERMANN, M., M. DIETZ & W. SCHORCH (2008): From a "Plattenbau" block of flats into a tower for bats. A report with hints for planning. 27 pp. Copies available from the EUROBATS Secretariat at eurobats@eurobats.org.
- BRIGGS, P. (2002): A study of bats in barn conversions in Hertfordshire in 2000. Hertfordshire Biological Records Centre, Hertford. Available on CD from HBRC, County Hall, Pegs Lane, Hertford SG13 8DN, UK.
- BRIGGS, P. (2004): Effect of barn conversion on bat roost sites in Hertfordshire, England. Mammalia: 68: 353-364.
- CATHERINE BICKMORE ASSOCIATES (2003): Review of work carried out on trunk road networks in Wales for bats. Catherine Bickmore Associates, London.
- DIETZ, C. (2001): Berücksichtigung des Fledermausschutzes bei der Sanierung von Natursteinbrücken und Wasserdurchlässen. Innenministerium Baden-Württemberg.
- DOWNS, N.C., V. BEATON, J. GUEST, J. POLANSKI, S.L. ROBINSON & P.A. RACEY (2003): The effects of illuminating the roost entrance on the emergence behaviour of *Pipistrellus pygmaeus*. Biological Conservation 111: 247-252.
- ELISONAS, J. (1932): Mūsu salies žinduoliai. Volume II. Svetimo ministerijos knygu leidimo komisija, Klaipėda.
- ENTWISTLE, A.C., P.A. RACEY & J.R. SPEAKMAN (1997): Roost selection by the brown long-eared bat *Plecotus auritus*. Journal of Applied Ecology 34: 399-408.
- ENTWISTLE, A.C., S. HARRIS, A.M. HUTSON, P.A. RACEY, A. WALSH, S.D. GIBSON, I. HEPBURN & J. JOHNSTON (2001): Habitat management for bats: A guide to land managers, landowners and their advisors. Joint Nature Conservation Committee, Peterborough, 47 pp.
- GRÉMILLET, X. (2006): Difficultés techniques d'isoler efficacement une colonie de parturition des sources diffuses d'intoxication mortelle (plomb et PCP): exemple d'une colonie de Grands Rhinolophes du Finistère. Actes des 10^{èmes} rencontres nationales « chauves-souris » de la SFPEM. Symbioses N.S. 14: 53-56.
- GRÉMILLET, X. & J. BOIREAU (2004): Intoxication mortelle par le plomb et par le fongicide PCP des juvéniles dans un gîte de parturition de Grands Rhinolophes, *Rhinolophus ferrumequinum*, dans le Finistère: difficultés de diagnostic et réalisation d'un gîte alternatif. Actes des 9^{èmes} rencontres nationales « chauves-souris » de la SFPEM. Symbioses 10: 59-61.
- HERNÁNDEZ, L.M., C. IBÁÑEZ, M.A. FERNÁNDEZ, A. GUILLÉN, M.J. GONZÁLEZ, & J.L. PÉREZ (1993): Organochlorine insecticide and PCB residues in two bat species from four localities in Spain. Bull Environ Contam Toxicol 50: 871-877.
- HUTSON, A.M. (1995): Conservation of bats in the management of ancient monuments. In: Managing ancient monuments: An integrated approach. Clwyd County Council, Clwyd: 71-78.
- IVANAUSKAS, T., N. LIKEVICIENĖ & S. MALDZIUNAITĖ (1964): Vadovas Lietuvos zinduoliams pazinti. Valstybine politines ir mokslines literatūros leidykla, Vilnius, 340 pp.
- JEFFERIES, D.J. (1976): Organochlorine insecticide residues in British bats and their significance. J. Zool. 166: 245-263.
- KELLEHER, C. & F. MARNELL (2006): Bat mitigation guidelines for Ireland. Irish Wildlife Manuals No. 25. National Parks and Wildlife Service, Dublin, 91 pp.
- KRAINER, K., C. DRESCHER & P. PRESETNIK (Eds.) (2007): Fledermausschutz im Alpen- und Adria-Raum 2003-2006. / Tutela dei Pipistrelli nell'area Alpina e Adriatica. / Varstvo dvošivk in netopirjev v regiji Alpe-Jadran. INTERREG IIIA Austria-Italia-Slovenia. ArgeNATURSCHUTZ, Klagenfurt, 80 pp.
- KURLAVICIUS P., D.H. PAUZA, V. MONSEVICIUS & S. GRUODIS (1991): Reti ir saugotini Lietuvos miskų gyvūnai. In: "Miskininko zinynas", Vilnius: Mokslas: 405-434.
- KURSKOV, A.N. (1981): Bats of Belarus. Nauka i Technika, Minsk, 136 pp. [Курсков А. Н. Рукокрылые Белоруссии. — Минск: Наука и техника, 1981. — 136 с.]
- KUZYAKIN, A.P. (1950): Bats. Sovetskaya Nauka, Moscow, 444 pp. [Кузякин А. П. Летучие мыши. — Москва: Советская наука, 1950. — 444 с.]
- LIMPENS, H., P.H.C. LINA & A. M. HUTSON (2000): Action Plan for the Conservation of the Pond Bat in Europe (*Myotis dasycneme*). Nature and Environment Series 108.
- LONGLEY, M. (2004): Greater horseshoe bat project 1998-2003. English Nature, Report No. 532.
- MASING, M.V. (1990): Bats of Estonia: current state of populations and ecological basics of conservation. Doctoral Thesis, Moscow, 233 pp. [Мазинг М. В. (1990): Рукокрылые Эстонии: современное состояние популяций и экологические основы охраны: Дис. канд.биол.наук — Тарту: ТГУ. — 233 с.]
- MITCHELL-JONES, A.J. (2004): Bat Mitigation Guidelines. English Nature. Peterborough, 74 pp.
- MITCHELL-JONES, A.J., G. AMORI, W. BOGDANOWICZ, B. KRYŠTUFEK, P.H.J. REIJNDERS, F. SPITZENBERGER, M. STUBBE, J.B.M. THISSEN, V. VOHRALÍK & J. ZIMA (Eds.) (1999): The atlas of European mammals. T & AD Poyser London, 484 pp. (Database EUNIS: <http://eunis.eea.eu.int>)
- MITCHELL-JONES, A.J., Z. BIHARI, M. MASING & L. RODRIGUES (2007): Protecting and managing underground sites for bats. EUROBATS Publication Series No. 2 (English version). UNEP/EUROBATS Secretariat, Bonn, Germany, 38 pp.



- (http://www.eurobats.org/publications/publication_series.htm)
- MITCHELL-JONES, A.J. & A.P. McLEISH (Eds.) (2004): *Bat Workers' Manual*, 3rd Edition. Joint Nature Conservation Committee, Peterborough, 178 pp.
- MOORE, N.P., S. JONES, A.M. HUTSON & D. GARTHWAITE (2003): Assessing the outcome of nature advice on bat colony management and mitigation works. English Nature, Report No. 517.
- NATIONALTRUST (2001): *Wildlife and Buildings. Technical guidance for architects, builders, building managers and others*. The National Trust, UK.
- PAINE, S. (1993): The effects of bat excreta on wall paintings. *The Conservator* 17: 3-10.
- PAINE, S. (unknown year): *Bats in churches. Guidelines for the assessment, identification and management of bat related damage to church contents*. Ancient Monuments laboratory, Conservation and Technology, English Heritage, London. (http://www.ribblevalley.gov.uk/downloads/file/232/bats_in_churches)
- PAUZA, D.H. (1985): *Kauno miesto ir jo apylinkiu šiksnosparniai*. Master Thesis. Vilnius, 131 pp.
- PAUZA, D.H. (1998): Distribution and status of Lithuanian Bats. *Proceedings of the Latvian Academy of Sciences Section B*, Vol. 52, No. 1-2: 594-596.
- PAUZA, D.H. & R. JUSKAITIS (1992): *Lietuvos Raudonoji Knyga. Žinduoliai*. Vilnius: Mokslas.
- PAUZA, D.H. & N. PAUZIENĖ (1983): *Inkilai šiksnosparniais. Mūsų gamta* 12, 13 pp.
- PAUZA, D.H. & N. PAUZIENĖ (1988): *Šiksnosparniai*. In: *Lietuvos fauna. Žinduoliai*. - Prūsaitė, J. (ed.) Mokslas, Vilnius: 43-91.
- PAUZA, D.H. & N. PAUZIENĖ (1996): Distribution, status and protection of Lithuanian bats. *Ekologija-Ecology-Экология*. Vilnius: Academia, No. 3: 44-65.
- PAUZA, D.H. & N. PAUZIENĖ (1998): *Bats of Lithuania: distribution, status and protection*. *Mammal review*. Vol. 28, No. 2: 53-67.
- PETERSONS, G. (1993): Distribution and present status of bats in Latvia. *Abstracts of the Second Baltic Theriological Conference*. Vilnius: 23.
- PRESETNIK, P. (2004): *Bat species and conservation issues in the castle Grad na Goričkem (NE Slovenia)*. *Mammalia* 68(4): 427-435.
- PRESETNIK, P., K. KLEMEN, M. ZAGMAJSTER, N. ZUPANČIČ, K. JAZBEC, U. ŽIBRAT, A. PETRINJAK & A. HUDOKLIN (2009): *Atlas netopirjev (Chiroptera) Slovenije / Atlas of bats (Chiroptera) of Slovenia*. Center za kartografijo favne in flore. Miklavž na Dravskem polju, 152 pp.
- PRESETNIK, P. & M. ZAGMAJSTER (2005): *Pod skupno streho z netopirji*. Informativna zloženska. Projekt Varstvo dvoživk in netopirjev v regiji Alpe-Jadran. INTERREG III A. Center za kartografijo favne in flore, Miklavž na Dravskem polju. (http://www.ckff.si/projekti/interreg/dokumenti/zgibankanetopirji_05.pdf)
- PRUSAITĖ, J. (1972): *Lietuvos žinduoliai*. Mintis, Vilnius.
- PYSARCZUK, S. & G. REITER (2008): *Bats and bridges in Austria*. *Abstracts of the XIth European Bat Research Symposium*, 18-22 August 2008, Cluj-Napoca: 121.
- RACEY, P.A. & S.M. SWIFT (1986): The residual effects of remedial timber treatments on bats. *Biological Conservation* 35: 205-214.
- RANSOME, R.D. & A.M. HUTSON (2000): *Action plan for the conservation of the Greater Horseshoe Bat in Europe (Rhinolophus ferrumequinum)*. Nature and Environment. N 109. Council of Europe Publishing, 53 pp.
- REITER, G. & A. ZAHN (2006): *Leitfaden zur Sanierung von Fledermausquartieren im Alpenraum / Bat roosts in the Alpine area: Guidelines for the renovation of buildings*. Co-ordination Centre for Bat Conservation and Research in Austria (KFFÖ) and Co-ordination Centre for Bat Conservation in South Bavaria + Department of Biology II, LMU Munich, 150 pp. (www.fledermausschutz.at/Sets/Literatur-Set.htm, section "download" => INTERREG III B Projekt)
- SARGENT, G. (1995): *The bats in churches project*. The Bat Conservation Trust, London.
- SCHOFIELD, H.W. (2008): *The lesser horseshoe bat conservation handbook*. The Vincent Wildlife Trust, Herefordshire, UK.
- SCHOFIELD, H. & F. BONTADINA (1999): *Habitat preferences of the lesser horseshoe bat, Rhinolophus hipposideros*. The Vincent Wildlife Trust Report.
- SHIEL, C. (1999): *Bridge usage by bats in County Leitrim and County Sligo*. The Heritage Council, Ireland. (http://www.heritagecouncil.ie/fileadmin/user_upload/Publications/Wildlife/Bridge_Usage_By_Bats_In_County_Leitrim_And_County_Sligo.pdf)
- SOCIÉTÉ FRANÇAISE POUR L'ÉTUDE ET LA PROTECTION DES MAMMIFÈRES (2002): *Les chauves-souris, hôtes des ponts*. SFEPM, France. (<http://www.sfepm.org/NuitChauveSouris/images2/Savoirplus/plaqponts.pdf>)
- SIMPSON & BROWN ARCHITECTS (Eds.) (1996): *The design and construction of bat boxes in houses*. Scottish Natural Heritage.
- SIMON, M., S. HÜTTENBÜGEL & J. SMIT-VIERGUTZ (2004): *Ökologie und Schutz von Fledermäusen in Dörfern und Städten / Ecology and conservation of bats in villages and towns*. Schriftenreihe für Landschaftspflege und Naturschutz, Heft 76 / 77. Bundesamt für Naturschutz, Bonn, 276 / 263 pp.
- SMITH, P.G. & P.A. RACEY (2002): *Habitat management for Natterer's bat Myotis nattereri*. Mammals Trust UK Publication. Peoples' Trust for Endangered Species, London.
- SWIFT, S.M. (1998): *Long-eared bats*. Poyser, London.
- TURNER, V.L. (2003): *Selection of foraging habitat by Daubenton's bat (Myotis daubentonii) and the two phonic forms of pipistrelle (Pipistrellus pipistrellus and P. pygmaeus) in a North Wales upland river catchment*. Countryside Council for Wales, Report No. 588.
- ZAGORODNIUK, I., L. GODLEVSKA, V. TYSHCHENKO, Y. PETRUSHENKO (2002): *Bats of Ukraine and adjacent countries: a guide for field investigations*. Kyiv, 108 pp. [Загороднюк, І., Годлевська, Л., Тищенко, В., Петрушенко, Я. (2002): *Кажани України та суміжних країн: керівництво для польових досліджень*. Київ. 108 с.] (<http://www.kazhan.org.ua/ukr/library/batguide.htm>)



Scientific and common names of European bats

Barbastella barbastellus – Western barbastelle bat
Barbastella leucomelas – Eastern barbastelle bat
Eptesicus bottae – Botta's serotine bat
Eptesicus nilssonii – Northern bat
Eptesicus serotinus – Common serotine
Hypsugo savii – Savi's pipistrelle
Miniopterus schreibersii – Schreibers' bat
Myotis alcaethoe – Alcaethoe whiskered bat
Myotis aurascens – Steppe whiskered bat
Myotis bechsteinii – Bechstein's bat
Myotis blythii – Lesser mouse-eared bat
Myotis brandtii – Brandt's bat
Myotis capaccinii – Long-fingered bat
Myotis dasycneme – Pond bat
Myotis daubentonii – Daubenton's bat
Myotis emarginatus – Geoffroy's bat
Myotis hajastanicus – Armenian whiskered bat
Myotis myotis – Greater mouse-eared bat
Myotis mystacinus – Whiskered bat
Myotis nattereri – Natterer's bat
Myotis nipalensis – Asiatic whiskered bat
Myotis punicus – Maghreb mouse-eared bat
Myotis schaubi – Schaub's bat

Nyctalus lasiopterus – Greater noctule
Nyctalus leisleri – Leisler's bat
Nyctalus noctula – Noctule bat
Otonycteris hemprichii – Hemprich's long-eared bat
Pipistrellus kuhlii – Kuhl's pipistrelle
Pipistrellus nathusii – Nathusius' pipistrelle
Pipistrellus pipistrellus – Common pipistrelle
Pipistrellus pygmaeus – Soprano pipistrelle
Plecotus auritus – Brown long-eared bat
Plecotus austriacus – Grey long-eared bat
Plecotus kolombatovici – Mediterranean long-eared bat
Plecotus macrotus – Mountain long-eared bat
Plecotus sardus – Sardinian long-eared bat
Rhinolophus blasii – Blasius' horseshoe bat
Rhinolophus euryale – Mediterranean horseshoe bat
Rhinolophus ferrumequinum – Greater horseshoe bat
Rhinolophus hipposideros – Lesser horseshoe bat
Rhinolophus mehelyi – Mehely's horseshoe bat
Rousettus aegyptiacus – Egyptian fruit bat
Tadarida teniotis – European free-tailed bat
Taphozous nudiventris – Naked-rumped tomb bat
Vespertilio murinus – Parti-coloured bat

Acknowledgements

The authors would like to thank the following EUROBATS delegates who had a significant active input into the Intersessional Working Group: Melanie Hardie (United Kingdom), Branko Karapandza (Republic of Serbia), Katie Parsons (BCT, United Kingdom), Dainius H. Pauza (Lithuania), Alison Rasey (BCT, United Kingdom), Dino Scaravelli (Italy), and Lubomira Vavrova (Slovakia).

Additional advice was also provided by: Stéphane Aulagnier (France), Andriy-Taras Bashta (Ukraine), Zoltán Bihari (Hungary), Marie-Jo Dubourg-Savage (SFEPM, France), Suren Gazaryan (Russian Federation), Lena Godlevska (Ukraine), Daniela Hamidović (Croatia), Jana Kristanc (Slovenia), Sandrine Lamotte (Belgium), Blanka Lehotska (Slovak Republic), Jean Matthews (United Kingdom), Branko Micevski (FYR Macedonia), Aleksandar Nastov (FYR Macedonia), Ioseb Natradze (Georgia), Marie Nedinge (Sweden), Guido Reiter (KFFÖ, Austria), Luísa Rodrigues (Portugal), Friederike Spitzenberger (Austria), Nikola Tvrčković (Croatia), and Marcel Uhrin (Slovakia).

Many thanks to The Center for Cartography of Fauna and Flora (Slovenia) for producing the maps used in this publication.

Photos are credited individually throughout the text. A number of extracts from *The Bat Mitigation Guidelines* (MITCHELL-JONES 2004) are incorporated into the text. These are also acknowledged as they appear.

Last but not least many thanks go to the national experts who completed the questionnaire: Ariana Koça (Albania), Mark

Kalashyan & Eduard Yavruyan (Armenia), Guido Reiter & Friederike Spitzenberger (Austria), Irina Rakhmatulina (Azerbaijan), Nico Verwimp (Belgium), Jasminko Mulaomerović (Bosnia and Herzegovina), Teodora Ivanova (Bulgaria), Nikola Tvrčković (Croatia), Eva Cepakova & Josef Chytil (Czech Republic), Hans J. Baagøe & Maj F. Munk (Denmark), Kaja Lotman & Matti Masing (Estonia), Eeva-Maria Kyheröinen & Torsten Stjernberg (Finland), Stéphane Aulagnier (France), Ioseb Natradze & Alexander Bukhnikashvili (Georgia), Peter Boye (Germany), Zoltán Bihari (Hungary), Ferdia Marnell & Kate McAney (Republic of Ireland), Stefania Biscardi & Dino Scaravelli (Italy), Gunars Petersons (Latvia), Michael Fasel & Silvio Hoch (Liechtenstein), Sigute Alisauskiene (Lithuania), Jacques Pir (Luxembourg), Aleksander Nastov, Svetozar Petkovski & Branko Micevski (FYR Macedonia), Peter H. C. Lina (the Netherlands), Per Ole Syvertsen, Øystein Størkersen & Kjell Isaksen (Norway), Wiesław Bogdanowicz (Poland), Luísa Rodrigues (Portugal), Dumitru Murariu (Romania), Suren V. Gazaryan (Russian Federation), Milan Paunović (Serbia including data from Montenegro), Peter Kanuch (Slovakia), Primož Presetnik & Jana Kristanc (Slovenia), Marie Nedinge & Johnny de Jong (Sweden), Coordination Centre for the Study and Protection of Bats (section east) (Switzerland), Lena Godlevska (Ukraine), and Rachel Harris (United Kingdom).



Annex 1: Questionnaire on dependence of bat species on overground roost types

Intersessional Working Group 5 - Protection of overground bat roosts

Resolution 4.9; 3a) states that:

Information on methods used to protect roost sites other than underground sites should be gathered by the Advisory Committee, with roost sites in buildings that are part of the cultural heritage as a priority.

QUESTIONNAIRE

Country:	_____
Completed by:	_____
Contact details:	_____
Date:	_____

General

- 1) Is there a national bat roost database for your country? _____ Y/N
- 2) If you know who holds this database, please give details here _____

- 3) Does it allow you to identify roost types? _____ Y/N

- 4) What type of overground roosts is used by which species of bats in your country?
Please indicate the dependence of individual species on specific roost types as High (H), Medium (M), Low (L), not important (-) or not known (?).

Species	Overground roost type						Other, please specify
	Church	Castle / fortification	House / block of flats	Barn / stable	Bridge	Tree	
<i>Rousettus aegyptiacus</i>							
<i>Taphozous nudiventris</i>							
<i>Rhinolophus blasii</i>							
<i>Rhinolophus euryale</i>							
<i>Rhinolophus ferrumequinum</i>							
<i>Rhinolophus hipposideros</i>							
<i>Rhinolophus mehelyi</i>							
<i>Barbastella barbastellus</i>							
<i>Barbastella leucomelas</i>							
<i>Eptesicus bottae</i>							
<i>Eptesicus nilsonii</i>							
<i>Eptesicus serotinus</i>							
<i>Hypsugo savii</i>							
<i>Myotis alcathoe</i>							
<i>Myotis aurascens</i>							
<i>Myotis bechsteinii</i>							
<i>Myotis blythii</i>							
<i>Myotis brandtii</i>							
<i>Myotis capaccinii</i>							
<i>Myotis dasycneme</i>							
<i>Myotis daubentonii</i>							
<i>Myotis emarginatus</i>							
<i>Myotis hajastanicus</i>							



4) (cont.) What type of overground roosts is used by which species of bats in your country?

Please indicate the dependence of individual species on specific roost types as High (H), Medium (M), Low (L), not important (-) or not known (?).

Species	Overground roost type						Other, please specify
	Church	Castle/fortification	House/block of flats	Barn/stable	Bridge	Tree	
<i>Myotis myotis</i>							
<i>Myotis mystacinus</i>							
<i>Myotis nattereri</i>							
<i>Myotis nipalensis</i>							
<i>Myotis cf. punicus</i>							
<i>Myotis schaubi</i>							
<i>Nyctalus lasiopterus</i>							
<i>Nyctalus leisleri</i>							
<i>Nyctalus noctula</i>							
<i>Otonycteris hemprichii</i>							
<i>Pipistrellus kuhlii</i>							
<i>Pipistrellus nathusii</i>							
<i>Pipistrellus pipistrellus</i>							
<i>Pipistrellus pygmaeus</i>							
<i>Plecotus alpinus</i>							
<i>Plecotus auritus</i>							
<i>Plecotus austriacus</i>							
<i>Plecotus kolombatovici</i>							
<i>Plecotus sardus</i>							
<i>Vespertilio murinus</i>							
<i>Miniopterus schreibersii</i>							
<i>Tadarida teniotis</i>							

Administrative and practical protection

- 5) Are bat roosts legally protected? _____ Y/N
If yes, please give details of the legislation (e.g. "Irish Wildlife Acts (1976 and 2000) protect all bat roosts from intentional disturbance or destruction"): _____
- 6) Are some roosts protected through state ownership or NGO ownership? _____ Y/N
- 7) Is there physical protection of overground roosts in your country? _____ Y/N
If yes, what forms of protection are used?
a) Grilling _____ Y/N
b) Fencing _____ Y/N
c) Blocking up _____ Y/N
d) Access restriction _____ Y/N
Other, please specify: _____

Interactions with built heritage

- 8) Can conservation of built heritage (e. g. churches, castles) conflict with bat conservation? _____ Y/N
- 9) What types of conflict arise?
a) Disturbance of bats by humans _____ Y/N
b) Disturbance of humans by bats (e.g. noise) _____ Y/N
c) Damage to property by bats _____ Y/N
d) Exclusion of bats from buildings/bridges / trees _____ Y/N
Other, please specify: _____
- 10) Please give an example, if you have one, of how such conflict has been successfully resolved.
- 11) Is it a legal requirement to carry out bat surveys before renovation / restoration works of buildings of cultural importance? _____ Y/N
- 12) Where bats are known to be present in a building of cultural importance due for renovation/restoration, are mitigation measures for the conservation of the bats legally required? _____ Y/N
If yes, what forms of mitigation are practised?
a) Timing of works to minimise disturbance _____ Y/N
b) Creation of limited access areas to protect bats _____ Y/N
c) Translocation of bats _____ Y/N
d) Provision of alternative roosts _____ Y/N
e) Exclusion of bats _____ Y/N
Other, please specify: _____
- 13) Is there information/education (e.g. advisory leaflets, training courses) available for owners of cultural heritage buildings/cultural heritage officials/architects etc. about bat conservation? _____ Y/N
If yes, please give brief details: _____

- 14) Please list relevant literature and/or web pages on conservation measures for bats in overground roosts.

Thank you for taking the time to complete this questionnaire!



Annex 2: Summary of questionnaire responses

Number of countries with estimated dependence of bat species on overground roost types.

Dependence: high (H), medium (M), low (L), not important (NI), not known (?); CH categories valid for Switzerland. Where respondents used a combination of categories (e.g. H/M) the higher dependence was taken; **bold** values highlight H+M dependence exceeding 4.

Species	Castle / fortification					Church					House / block of flats (CH buildings)					Barn / stable					Bridge (CH bridge / rock)					Tree				
	H	M	L	NI	?	H	M	L	NI	?	H	M	L	NI	?	H	M	L	NI	?	H	M	L	NI	?	H	M	L	NI	?
<i>Barbastella barbastellus</i>	2	5	2	8	12	2	5	10	12	3	3	6	5	12	4	6	5	14		2	9	18	10	2	3	1	13			
<i>Barbastella leucomelas</i>			1	1	2			1	1	2			1	3				4				1	3			1	1	2		
<i>Eptesicus bottae</i>		1	1		1			1	2	1				2				3				1	2	1	1			1		
<i>Eptesicus nilssonii</i>	2	3	4	6	7	1	5	5	8	4	8	6	1	2	6	3	1	7	12		11	12	2	2	3	5	11			
<i>Eptesicus serotinus</i>	4	4	3	5	15	9	5	5	3	9	17	3	2	9	2	3	4	6	16	2	2	11	16	1	5	8	17			
<i>Hypsugo savii</i>		3	2	5	9		3	7	9		3	4	2	3	7	1	1	4	13	1	1	5	12	1		6	12			
<i>Miniopterus schreibersii</i>	1	1	3	7	6	1	1	3	7	6		3	9	6		1	8	9		9	9					10	8			
<i>Myotis alcaethoe</i>				6	2			6	1			1	3	4			5	3			5	3	2			3	3			
<i>Myotis aurascens</i>				4	1			5	1			2	2	1		3	1	1	1	3						4	1			
<i>Myotis bechsteinii</i>		1	4	8	13		1	13	12		2	3	7	14		1	11	14		1	10	15	17		1		8			
<i>Myotis blythii</i>	3	4	3	5	6	7	4	3	4	3	3	1	4	5	8	1	1	1	8	10	1	2	1	6	11		1	9	11	
<i>Myotis brandtii</i>	4	1		7	15		2	3	8	14	7	8		4	9	2	1	5	19	1	9	17	9	2	1	2	13			
<i>Myotis capaccinii</i>			2	5	5			1	6	5			1	6	5			1	5	6	1					4	7		5	7
<i>Myotis dasycneme</i>	3	2	1	3	8	5	2		4	6	8	2	1	1	5	2		5	10			5	12	1	4	2	1	9		
<i>Myotis daubentonii</i>	6	3	4	4	15	1	1	4	11	15	3	6	6	4	13	5	2	9	16	6	6	7	2	11	21	2		1	8	
<i>Myotis emarginatus</i>	6	7	1	3	8	8	5	2	4	6	4	5	4	3	9	3	2	4	5	11	1	1	9	14			8	17		
<i>Myotis hajastanicus</i>				1	1			1	1				1	1				1	1		1	1				1	1		1	1
<i>Myotis myotis</i>	5	5	2	2	7	11	4		2	4	4	3	3	5	6	3		9	9	2	3	6	10	1		8	12			
<i>Myotis mystacinus</i>	5	2	4	6	18	2	2	7	7	17	12	7	1	3	12	4	2	3	6	20	1	2	1	9	22	8	3	2	3	19
<i>Myotis nattereri</i>	4	4	3	3	21	2	1	5	7	20	7	6	1	3	18	2	4	2	3	24	1	3	2	8	21	10	1	1	22	
<i>Myotis nipalensis</i>				1				1					1					1			1							1		
<i>Myotis punicus</i>		1		2				3				1	2				1	2			1	2					3			
<i>Myotis schaubi</i>				1	1			1	1				1	1				1	1		1	1				1	1		1	1
<i>Nyctalus lasiopterus</i>		1		5	5			1	6	4		1	1	4	5		1	5	5			5	6	8				4		4
<i>Nyctalus leisleri</i>		2		11	18			1	14	16	1	4	5	6	15		1	1	10	19		2	10	19	18		1		12	
<i>Nyctalus noctula</i>	1	1	1	12	18	3	1	5	9	15	9	5	3	3	13	1		1	10	21	1	1	3	8	20	23	2		8	
<i>Otonycteris hemprichii</i>				1				1					1					1			1							1		1
<i>Pipistrellus kuhlii</i>	2	4	1	5	9	3	3	3	6	6	16	1		1	3	4	1	3	4	9	2	2	6	11	2	1	1	5	12	
<i>Pipistrellus nathusii</i>		2	4	10	16	2	3	4	11	12	13	6	3	2	8	3	1	3	6	19	2	1	10	19	18	3	1	1	9	
<i>Pipistrellus pipistrellus</i>	5	4	4	4	18	8	3	5	6	13	20	5		2	8	5	3	5	4	18	3	2	4	7	19	8	3	3	2	19

Summary of questionnaire responses (cont.)

Species	Castle / fortification					Church					House / block of flats (CH buildings)					Barn / stable					Bridge (CH bridge / rock)					Tree				
	H	M	L	NI	?	H	M	L	NI	?	H	M	L	NI	?	H	M	L	NI	?	H	M	L	NI	?	H	M	L	NI	?
<i>Pipistrellus pygmaeus</i>	3	2		7	15	4	1	3	9	10	10	5		3	9	3	1	2	2	19	2	1		7	17	9	4	1	2	11
<i>Plecotus auritus</i>	8	5	5	1	15	15	6	5	4	4	12	9	3	1	9	2	6	6	3	17	2	1	9	22	11	5	2	1	15	
<i>Plecotus austriacus</i>	6	2	3	2	13	11	3	4	2	6	9	3	3		11	3	1	3	4	15	2		7	17	2	2		5	17	
<i>Plecotus kolombatovici</i>				1	1	1			1				1					1	1					1	1				1	1
<i>Plecotus macrobullaris</i>		2		3	5	3	4	1	1	1	3	3		1	3	1		1	4	4			5	5	2	1	3	4		
<i>Plecotus sardus</i>				2				2		1			1					2			2								2	
<i>Rhinolophus blasii</i>		1	2	3	4		1	1	3	5			1	3	6			1	3	6			3	7				3	7	
<i>Rhinolophus euryale</i>	1		4	8	5	2		2	7	7			3	7	8			2	8	8			1	6	11				9	9
<i>Rhinolophus ferrumequinum</i>	6	5	5	1	8	7	4	4	3	7	3	8	5	2	7	2	3	6	5	9	2	1	2	6	14				13	12
<i>Rhinolophus hipposideros</i>	7	6	4	2	6	6	5	4	6	4	8	7	3	1	6	4	3	8	4	6	1	2	3	7	12		1	13	11	
<i>Rhinolophus mehelyi</i>		1		6	5		1	1	5	5			5	7				5	7				5	7				7	5	
<i>Rousettus aegyptiacus</i>				1				1					1					1					1						1	
<i>Tadarida teniotis</i>		1		3	10		1	2	5	6	1	1	1	1	10			4	10	1		1	3	9	1			4	9	
<i>Taphozous nudiventris</i>				1				1					1					1					1						1	
<i>Vespertilio murinus</i>	1	1	3	7	16	1	1	6	8	12	15	6	1	1	5	3		2	5	18		1	1	7	19	2	1	2	4	19



Annex 3: EUROBATS Resolution 5.7

EUROBATS.MoP5.Record.Annex10

5th Session of the Meeting of Parties
Ljubljana, Slovenia, 4 – 6 September 2006

Resolution No. 5.7



Guidelines for the Protection of Overground Roosts, with particular
reference to roosts in buildings of cultural heritage importance

The Meeting of the Parties to the Agreement on the Conservation of Populations of European Bats (hereafter “the Agreement”),

Recalling Resolution No. 4.9, Element 3 (a) on the protection of roost sites other than underground sites;

Recognising the importance of buildings as roost site for many species of bats, both for hibernation and breeding in different parts of their ranges;

Further recognising that buildings, which contain bat roosts, may also in themselves be of built heritage importance and further recognizing that protection and restoration works may be required for such structures;

Urges the Advisory Committee to complete and publish the guidance document (now in draft form);

Urges Parties and Non-Party Range States to:

1. Establish national databases of important overground roosts;
2. Ensure that the important overground roosts they have identified are fully protected by law and where appropriate, are physically protected against unauthorised entry;
3. Establish a working relationship between the relevant cultural and natural heritage agencies, including, where appropriate, linkage between databases of bat roosts and databases of cultural heritage buildings;
4. Include in future national reports a summary of these interactions;
5. Prepare guidelines for custodians of historical buildings, on the protection of bat roosts;
6. Develop schemes, which encourage the maintenance of bat roosts in buildings of cultural heritage, including, if appropriate, grants to ensure the maintenance of bat roosts during renovation/restoration;
7. Encourage architects and engineers to incorporate a natural heritage element into their training programmes;
8. Ensure that overground roosts are managed in accordance with national nature conservation legislation and taking note of any guidelines adopted by the EUROBATS Agreement.

Annex 4: Summary of good practise for the protection of overground roosts in buildings of cultural heritage importance

Many European bat species have shown significant declines in recent decades and several species are highly endangered. In almost all European countries, bats are protected by local, national and/or international legislation or Agreements (*e.g.* Habitats Directive, Bonn Convention). Achieving and maintaining favourable conservation status for bats requires that their breeding and resting sites – roosts – are also protected (see Habitats Directive).

In Europe, a high percentage of bat species roost for at least part of each year in buildings. Buildings of cultural heritage importance are often of particular importance for bats. These structures may be protected in their own right. This report provides practical advice on how to manage bat roosts in buildings including those of cultural heritage importance.

Bats and buildings can come into conflict in two ways:

1. When bats are causing damage (see page 28 ff for details)

The priority must be to maintain the bats roost as intact as possible. Non-invasive mitigation measures should have priority over more invasive actions. Before acting seek advice from the local bat group or statutory nature conservation organisation (SNCO).

Examples of easy and inexpensive actions:

- Remove droppings;
- Move or cover important objects;
- Install deflector boards.

Examples of more complex actions which usually require special licences and official SNCO approval:

- Relocation of roost access;
- In extreme situations, relocation of bats to a nearby replacement roost may be the only option.

2. When renovation or maintenance works are required (see chapter 6 for case studies)

Good planning is essential. If the works are planned well in advance there will be time to gather all necessary information about the bats using the building. This will allow works to be scheduled to minimise the impact on bats, thereby reducing the need for costly mitigation measures and work stoppages. In EU countries disturbance of bat roosts may require a derogation licence or, in Natura 2000 sites, may also require an environmental impact assessment. It is wise, therefore, to seek early advice from the SNCO.

- Establish which bat species are present, where they roost, their access points and seasonality of use. This information may already be known (*e.g.* by a local bat group). If not, a full bat survey, ideally covering all seasons, should be undertaken by a bat expert.
- If bats are present in the building, it is wise to include the bat expert into the engineering team.
- Time the works to avoid disturbing the bats.
- Do not block access points or damage roost.
- Ensure materials to be used are bat friendly.
- Enhance bat roosting potential where feasible.
- Monitor effectiveness of conservations measures.